

Sportfish Management in Ponds

Gary J. Burtle, Extension Fisheries Specialist

P. O. Box 748

Tifton, GA 31793-0748

Telephone: 229-386-3364; E-mail: fish@tifton.uga.edu

Contents

Stocking

Fertilization and Liming

Supplemental Feeding

Fishing Methods

Sportfish ponds and small impoundments provide significant opportunity to enjoy the outdoors and enhance property value. Properly managed ponds provide an abundant supply of fish for recreation and nutrition. The presence of water attracts wildlife of a variety not possible with dry land. Livestock can utilize ponds as a source of drinking water. Ponds can be adapted to provide a source of water for fire protection in rural areas. The calming effects of a body of water provides added value to real estate in urban and rural settings. These are some reasons why millions of ponds have been constructed in the United States and why pond construction and renovation continues.

Fish Stocking

Stock fish according to your goals for fish production and fishing preferences. Balance can be managed for combinations of largemouth bass, sunfish, and channel catfish. Other species can be added as forage for the bass, but adding fish that compete with the bass or sunfish can change the balance in a negative way. The initial stocking rates are also set by your fertilization plans. More fish can be stocked when a routine fertilization schedule is followed than when a pond is not fertilized.

Largemouth bass are carnivorous fish that usually are not trained to accept a commercial feed. A forage species must be stocked with the bass to provide the necessary food for them to grow. In most cases the forage species are bluegill sunfish and red ear sunfish. However, fathead minnows and threadfin shad are sometimes stocked to supplement the sunfish forage base. In some cases hatchery-trained bass will eat pellets of a high-protein feed. Automatic feeders are used to dispense feed at regular intervals so that bass are not allowed to revert back to their forage feeding instincts.

Sunfish are also enjoyed as food and sport by many fishermen and complement the largemouth bass in a pond community. The combination of bluegill sunfish and red ear sunfish allows for diversity in the pond by recognizing that the red ear sunfish eat crustaceans and mollusks to a greater extent than the bluegill sunfish do. Snail control can be accomplished when the proper number of red ear sunfish are stocked and maintained in a pond. Sunfish are prolific spawners that reach spawning age within a few months after hatching. Without a carnivore, like the largemouth bass, sunfish would soon overpopulate the pond and become stunted.

Channel catfish are a fast-growing source of food and fishing enjoyment for the fisherman. Enough catfish are stocked to meet the fishing plans of the pond owner then restocked when most are caught. Small catfish may be eaten by largemouth bass and sunfish may eat catfish eggs if spawning occurs. However, catfish spawning may result in pond overcrowding if fishermen do not harvest enough catfish. Accumulations of large catfish or small catfish can cause serious pond imbalance. A commitment to catfish harvesting must be made before the channel catfish are stocked.

Sources of Fish

Private hatcheries are the major source of fish for stocking private ponds in Georgia. Over the years, hatcheries have been developed throughout the state. These hatcheries provide fish to distributors for more complete coverage of the state. Delivery is available if the pond owner cannot transport fish from the hatchery. Small quantities of fingerling size fish can be transported in plastic bags under an atmosphere of oxygen. Some private hatcheries, usually from out-of-state, arrange to meet pond owners at a central location for fish pick-up.

The health and quality of fish should be excellent when considering the source for stocking. Check the fish quality by talking to past customers or visiting the

hatchery before fish are purchased. Dead fish in holding tanks and fish with sores or ragged fins are signs of problems. When fish are delivered, check the fish before they are unloaded. Fish should be tempered or acclimated to the receiving pond water before unloading in order to avoid adding stress to the transport experience. Cool weather and cool water reduce the possibilities of transport stress of fish.

Special strains of largemouth bass or bluegill sunfish may be obtained from private hatcheries. The Florida strain of largemouth bass (*Micropterus salmoides floridanus*) and its hybrid with the Northern largemouth (*Micropterus salmoides salmoides*) grow faster and have significantly better survival for the first three years than the Northern largemouth bass. Angling success is not different among these strains of bass. Bluegill (*Lepomis macrochirus macrochirus*) and coppernose bluegill (*Lepomis macrochirus purpureus*) may be distinct strains and the coppernose has been reported to grow faster than wild bluegill. The coppernose bluegill has a distinct copper colored band across the front of its head. Coppernose bluegill should not be confused with hybrid bream that are interspecific crosses.

Winter stocking rainbow trout has become a common practice in Georgia ponds. A large fingerling (six inches) should be stocked in November or December for harvest during February to March. Catchable trout may be stocked in December for fishing till the end of March. Rainbow trout may be obtained from private hatcheries in Georgia or states to the north.

Fish for sportfish ponds should not be obtained from public lakes or streams. Wild fish may not be as adaptable to small static water ponds as domesticated hatchery fish. Disease organisms may be found on any fish, but are likely to be carried by wild fish obtained from uncontrolled waters. Transfer of diseased fish from one body of water to another is not legal in Georgia.

Stocking Rates

Bass-Bream or Bass-Bream-Catfish combination are two popular stocking choices (Table 1). These fish combinations were made popular after research during the 1940's and '50's showed that a balance between largemouth bass and bream (bluegill and redear sunfish) could produce good fishing. Some differences in the stocking numbers are reported for different locations, but this basic approach to sportfish pond stocking is widely accepted. The sunfish are stocked first in the fall to early spring so that they can spawn before the bass are stocked in early summer.

Catfish are usually stocked in the fall or winter when they are most available. Triploid grass carp are utilized to control aquatic plant growth in ponds and should be stocked in new ponds and ever five or six years thereafter.

Table 1. Stocking rates for sportfish ponds in Georgia.		
Fish Species	Number per Acre of Fertilized Pond	Number per Acre of Unfertilized Pond
Bream (70-80% bluegill and 20-30% redear sunfish)	500 to 1000	500
Largemouth Bass	100	50
Channel Catfish (optional)	100 to 500	100
Rainbow Trout (winter only)	100 to 300	100
Sterile Grass Carp	5 to 10	5

The higher rates for bream stocking are used when trophy largemouth bass are preferred. Additional forage will be available for more rapid bass growth. When the Florida largemouth is stocked, 1,000 bream should be stocked. When catfish are stocked at the highest rates, a regular feeding program should be utilized. Rainbow trout stocked at the higher rate should be fed a high quality trout or salmon feed. Rainbow trout stocked at small sized with large bass may be eaten as forage.

Other Stocking Options

Fish production goals sometimes include hybrid bream, hybrid striped bass, crappie, bullhead catfish, or forage fish. Each of these fish would change the balance of a bass-bream population. Some can enhance the bass-bream balance but most have undesirable consequences. For that reason, these fish are usually stocked alone, in small ponds, or in ponds over 10 acres in size that may be large enough to allow them to coexist with largemouth bass and sunfish.

Hybrid bream need ponds with no outflow and no possibility of contamination with other sunfish. A common hybrid sunfish is produced by crossing the bluegill sunfish with the green sunfish. Other hybrids are more complex crosses among species. The growth potential for the hybrid depends on the genetic background of each parent strain or species. Most hybrid sunfish are reported to grow faster than the bluegill, however not all may be from domesticated strains selected for fast growth. Therefore, investigate the source of hybrid sunfish carefully and obtain referrals from previous hatchery customers. Hybrid sunfish must be harvested on a regular schedule in order to make room for the potential of growing trophy size bream. Some individuals in the hybrid population will always reproduce, although not as productively as the bluegill or green sunfish parents. Reproduction among the hybrid bream results in reversion to the adult species through a series of intergrades (progeny with varying degrees of the characteristics of the original species used to create the hybrid). This process continues for only a few generations before a population resembling green sunfish develops. Active harvest of hybrid sunfish can postpone development of a population of crowded intergrades. Reproduction can be controlled with some success by stocking largemouth bass with hybrid bream and allowing the pond to remain bass overcrowded.

Hybrid striped bass are available for pond stocking, but require a steady supply of forage fish or feed to be provided. When a good supply of food is available, hybrid striped bass will grow to two pounds in two years. The cross between striped bass and white bass is most common. Bluegill sunfish are not a preferred forage species for hybrid striped bass and balance fish populations do not occur when these species are stocked together. Fathead minnows or threadfin shad are better choices as forage for hybrid striped bass.

Crappie, or speckled perch, are a favorite sportfish in Georgia. However, crappie are difficult to balance with other fish species. Large ponds of 10 acres or more may be more suitable for crappie-bass-bream populations. Crappie spawning is inhibited by high fish densities. Crappie like relatively clear water in order to see their prey and may not grow well in fertilized ponds with limited visibility. Largemouth bass spawns are eaten by young crappie that spawn earlier and at cooler water temperatures. Many attempts to produce crappie with bass-bream populations end up as overcrowded crappie-bluegill ponds.

Bullheads are traditional catfish species in Georgia. The brown bullhead (*Ameiurus nebulosus*) and yellow bullhead or butter cat (*Ameiurus natalis*) are

preferred for their delicate meat. Bullheads rarely grow larger than eight or nine inches in length although records indicate they can exceed 18 inches. In bass-bream populations, bullheads compete with bream for food and may reduce spawning success by eating eggs out of the nest. Bullheads can be stocked in small ponds by themselves for best fish production. Feed regularly to improve bullhead growth.

Forage fish may be stocked with bass-bream populations to obtain certain management goals. Ponds intended for production of trophy bass need extra forage and special management. When the forage base is bream, aggressive largemouth bass strains may be limited by a shortage of food. Bass in excess of 10 pounds may be produced with additional forage. Fathead minnows should be purchased at least twice a year as a forage supplement. Threadfin shad may establish reproducing populations in some ponds. Obtain threadfin shad from knowledgeable sources who can supply a pure population of threadfin without gizzard shad that look similar to the threadfin when small.

Channel catfish are often stocked by themselves in small ponds that are too small to balance bass-bream populations. When ponds are unfertilized, stock 100 to 200 channel catfish per acre. Fertilization allows an increase to 300 to 500 catfish per acre. Regular feeding is needed to allow stocking between 500 and 2,000 channel catfish per acre. Catfish should be harvested four to six months after stocking. Aggressive harvests of catfish should be used to remove most of the initial stocking before the end of the second year. Channel catfish may begin to spawn during the third or fourth year and cause overcrowding in the pond. Once thought to be difficult to spawn in ponds, channel catfish are now likely to spawn with a minimum of structure beside a board or piling, under a cement block, in a tire, or between tree roots.

Rainbow trout can be produced in earthen ponds as long as water temperatures are cool. At higher elevations in North Georgia, ponds may remain below 70° F for the entire year. However, in South Georgia, rainbow trout can be raised between November and March. Best growth is achieved when fewer than 1,000 are stocked per acre. As many as 3,000 rainbow trout per acre are stocked for heavy fishing pressure.

Fertilization

When the pond owner wants more fish to catch, fertilization allows a greater

production capacity for the pond. Unfertilized ponds may produce less than 100 pounds of fish per acre. If that is your goal, fertilization may not appeal to you. Fertilization will increase the amount of plant growth in a pond and should cause a bloom of phytoplankton (microscopic algae). Normal blooms allow visibility of 12 to 18 inches into the pond water, using a Secchi disc or light colored object held downward into the water. Phytoplankton and bacteria grow on the fertilizer nutrients and, in turn, feed zooplankton (microscopic animals). Zooplankton is a food for young fish and filter feeding fish that provide the forage base for largemouth bass. In this way, inorganic nutrients in fertilizer are converted into extra fish mass.

Fertilizers are available in a variety of forms and each pond owner may develop their own preference. Granular fertilizers have been used to increase fish mass since the concept of bass-bream ponds was first developed. Liquid fertilizers are relatively affordable and may be easier to apply than granules. Powder fertilizers are more expensive, but do not require dilution before application. Timed or slow release fertilizers are most expensive but allow season-long fertility in one application.

Fertilizer rates for fish ponds were originally based on the rate of 8 pounds of soluble phosphate per surface acre (Table 2). Balanced nitrate and phosphate formulations were used to develop that recommendation. When liquid fertilizers became popular, a ration of 1:3, nitrate to phosphate, was utilized in order to maintain the liquid form of the fertilizer. Recently, nitrate to phosphate ratios have approached 1:5 in powdered fertilizers and slow release fertilizers. Fertilizer results will vary from pond to pond because of differences in pond water alkalinity and soil chemistry. High alkalinity water reduces the solubility of powdered fertilizer. Soil chemistry determines the amount of phosphate that can remain dissolved in the pond water column. This relationship will be presented in the section on liming.

Fertilizer application requires that the fertilizer be kept from contact with the pond soil. Platforms or water soluble fertilizers are used to dissolve the nutrients into the water column where algae can utilize them. Phosphate will bind to clay in soil if allowed to fall to the bottom of the pond.

Granular fertilizers are usually placed on a platform where water currents can dissolve the nutrients into the pond water column. A platform can be constructed from decay resistant lumber that is six feet square and sets about 18 inches below

the water surface. Some platforms may have a walkway to allow easier access. An alternative method utilizes the plastic bag that fertilizers usually are shipped in. By laying the bag in shallow water and cutting an X across the upper flat side of the bag, nutrients will dissolve into the water column over time. Bags are staked to the pond bottom to prevent them from floating away when all of the fertilizer is dissolved. Slow release fertilizers may be sold with a plastic tray included. The tray is staked to the pond bottom and the fertilizer is placed inside.

Fertilization is begun when water temperature stabilizes above 60° F, possibly in late February and no later than April. Application may be repeated at two week intervals until a phytoplankton bloom develops. Applications are then based on the density of the bloom and are applied about once each month until October whenever the visibility exceeds 18 inches. It is common to make 10 or 12 fertilizer applications per year. Liquid fertilizer application follows the same pattern. Powders also follow that pattern, unless they are utilized with slow release fertilizer. Before placement of slow release fertilizer on a platform or tray, apply one application of powdered fertilizer to boost the bloom. In static water, the slow release fertilizer will provide bloom sustaining nutrients for the entire season.

Table 2. Examples of fertilizers used in sportfish ponds.		
Form	Composition N - P - K (%)	Amount per Application*
Granular	20-20-5	20-40 lb/A
Granular	10-10-10	40-80 lb/A
Granular	8-8-8	50-100 lb/A
Granular (Foating Bucket)	18-46-0	4 - 10 pounds/A
Liquid	10-34-0	1 gallon/A
Liquid	10-30-0	1 gallon/A
Liquid	13-37-0	1 gallon/A
Powder	12-48-8	4 - 10 pounds/A
Powder	10-52-4	4-8 pounds/A
Coated Granule (Slow Release)	10-50-0	25-40 pounds/A Annually

*Water with high alkalinity may require over 4.0 pounds of phosphate per acre.

Liming

Ponds are limed in order to change the soil pH from acid to slightly basic. Most soils in Georgia require some liming, however soils with limestone or marl deposits will probably not need to be limed for fish pond management purposes. Also, ponds that are filled using well water from limestone aquifers need little or no liming. Changing the acidity of soils allows them to release phosphorus more easily so that plants can utilize this nutrient for growth. Calcium carbonate in lime buffers the pH in the pond water so that changes during the day may not be great enough to cause distress to the fish.

The amount of lime needed per acre can be determined by analysis of a soil or water sample. Soil pH can be checked by submitting a sample to your county Extension agent. Water is used to determine lime requirement by checking the pond water alkalinity or total hardness. Water pH may not be a good indicator of lime requirement since the pH changes during the day due to the release and utilization of carbon dioxide by aquatic organisms. In some cases pond water pH may indicate the presence of dissolved organic acids like tannic acid. Certain ponds may have deposits of peat or decaying leaves that lower pH to levels that are too acid to support fish life.

When pond water alkalinity is above 20 mg calcium carbonate per liter, additional lime is not recommended. Enough phosphorus is released from the pond soil at that alkalinity to allow phytoplankton to grow, supplying the basis for the pond food chain. Phytoplankton blooms are stimulated by the fertilization schedules mentioned previously. Water hardness should also be above 20 mg calcium carbonate per liter. Hardness is a measure of elements including calcium, magnesium, manganese, and iron in the water (elements with two positive charges when dissolved in water). Hardness and alkalinity may each be utilized to determine lime requirement. Pond water hardness has been correlated to the cation exchange capacity of pond soils which is an indicator of lime requirement.

Lime requirement should be checked periodically in order to allow effective pond fertilization. Rainfall and other sources of water that flow through a pond cause water hardness and alkalinity to decrease over time. Ponds constructed across streams may not be able to retain enough alkalinity or fertilizer to allow management of phytoplankton blooms. Most ponds are limed but once every four

years.

Agricultural lime is the most common source of lime. Agricultural lime is a combination of calcium and magnesium carbonates. It is recommended that about 2 mg magnesium per liter of pond water should be available in fish ponds. Several types or grades of agricultural lime are available. The coarser types may be less soluble and lime from some sources may be hard to dissolve due to their mineral composition, but a lime that passes through 400 mesh screen is best for ponds. Lime takes about three months to dissolve and should be applied that far ahead of the spring fertilization period. Liquid lime (a suspension of fine particles of agricultural limestone) is useful because it dissolves quickly and can be easier to apply than bulk granular limestone.

Hydrated lime (builders lime or calcium hydroxide) can be applied to ponds to satisfy the lime requirement. More soluble and finely powdered, hydrated lime may increase pond water pH rapidly, causing a fish kill. It is best to apply less than 100 pounds of hydrated lime per surface acre at weekly intervals until the lime requirement is met. Hydrated lime application may need to be repeated every six to eight weeks in order to prevent a decline in water hardness.

Liming ponds allows use of algicides that contain copper. Since copper is more toxic when pond water alkalinity is low, lime can be used to increase alkalinity and reduce the toxicity of copper to fish. Trout, koi carp, and blue catfish have a lower tolerance to copper than do bream, bass, grass carp, or channel catfish. It is important to avoid using copper unless the alkalinity of pond water is higher than 40 mg per liter.

Feeding

When channel catfish are stocked in ponds, supplemental feeding is very important. Stocking rates of less than 100 catfish per acre can be sustained for a few years without feeding, but feeding should begin after the third year. Channel catfish spawn readily in ponds using any shelter they can find or digging out under the pond bank. Over the years, domestication of catfish has overcome their reluctance to spawn in static water ponds. So, catfish can restock themselves and overstocking can occur with uncontrolled catfish spawning. Supplying adequate feed for the catfish until they can be caught is necessary to prevent stress in the fish and disease infection following stress. In order to get rapid catfish growth, feed each day all they will eat in a 15 to 20 minute period. Avoid feeding during cloudy

weather when oxygen depletion is a danger. Discontinue feeding during very cool weather when water temperatures are below 65° F.

Supplemental feeding can improve bream growth under certain conditions but do not think that all bass-bream ponds must be fed. Bluegill sunfish have been grown to over one pound in size by using supplemental feeding. A high quality fish feed of 1/8 to 1/4 inch diameter should be used to feed bream. The feed should also contain between 32 and 36% protein. As with catfish, feed what the fish will consume in less than 15 to 20 minutes. However, bream are not fed in the sense that commercial, high density fish farms are, but special formulations are used as a supplemental ration. Automatic feeders can be utilized to make the feeding more successful by insuring regular feeding and allowing several feedings per day. Hybrid sunfish will also benefit from supplemental feed. However, stunted bream populations cannot be improved by using supplemental feeding. Other methods for correcting bream overcrowding will be explained later.

Feed amounts must be adjusted seasonally. Fish eat more in warm weather than in cold. It is not necessary to feed catfish or bream when water temperatures cool to less than 70° F. Also, in the hottest part of summer, feed amount should be reduced to account for slower feed consumption and the risk of oxygen depletion. Some feed reduction during changes in weather may also be advisable. Feed quantity or frequency can be reduced for a few days as a low pressure front passes by. Feed should be reduced or withheld after a few days of cloudy weather. When feeding more than 30 pounds of feed per acre per day, aeration should be available to protect from eventual oxygen depletion. In most bass-bream ponds, 10 pounds of feed per acre per day is adequate to support the fish. It is important to remember that oxygen depletion can occur even when ponds are not fed.

Fishing Methods

The first step to good pond management is setting goals for the type of fishing that you want. Once a goal is established, rules for catching, releasing, feeding or fertilizing can be set.

Fishing should not begin until June of the year following bream stocking and largemouth bass should not be removed until the second year after a successful spawn has been observed. This rule applies to both any bass-bream pond because the biology of both species requires managed harvest to prevent overcrowding of bream or over-fishing of bass. Several points must be considered:

1. Catching bream as soon as they become mature (3-4 months into first growing season) starts the bream harvest and helps the bass control bream reproduction.
2. Preserving bass until after their first spawn allows the bass to help manage bream numbers and provides bass replacements for the bass that are caught.
3. Largemouth bass diminish in numbers even without fishing and male largemouth bass seldom reach five years of age.
4. Removal of intermediate sized bream (4-6 inches) and bass (14 to 18 inches) helps develop a population of trophy sized fish by removing competition.

Table 3. Catch rates for bass-bream ponds.		
Species	Unfertilized Pond	Fertilized Pond
Bream (first year)	40 pounds (120 fish)	80 pounds (320 fish)
Bream (later years)	40 pounds (120 fish)	150 pound (600 fish)
Largemouth bass (second year)	10 pounds (8 fish)	20 pounds (15 fish)
Largemouth bass (later years)	10 pounds (5 to 8 fish)	35 pounds (15 to 20 fish)

Bream serve as the forage base for largemouth bass and must be managed to provide enough food for bass growth but not allowed to overpopulate and stunt their own growth. Bream must be removed on a regular basis from the pond, either by the bass or by anglers. Set goals for bream fishing based on Table 3 and monitor the size of bream caught. It has been said, “Never return bream to the pond, regardless of size.” In overcrowded bream ponds, a bluegill may be only three inches long, but fully mature. Removal of intermediate sized bream, between 4 and 6 inches long, will encourage growth of the larger bream to trophy size. Due to the relatively slow growth of bluegill sunfish, plan to wait four years until trophy bluegill, above one pound, are caught. Since bluegill seldom exceed six years of age, remove bluegill when they are caught since you may not get a chance to watch the big ones grow any larger. Large ponds and lakes may develop larger bluegill because more food is available and more space is available to allow the fish to escape capture by bass or anglers while they grow to trophy size.

Largemouth bass management depends on control of bass harvest. It takes a commitment to record keeping and catch limits in order to maintain a pond with bass that you may want to hang on your wall (10 pounds or larger). Use the catch limits in Table 3. Catch bass and remove when they reach 14 to 18 inches. Also, check for signs of bass overcrowding, then remove any stunted 10 to 12 inch bass. Bass stunting from overcrowding is an indicator of a pond that is under fished. Ponds with little fishing pressure will not produce the weights and numbers of fish that are listed in this publication. Abstaining from fish harvest will allow the pond to revert to a wild condition with low fish numbers of relatively small size. If a fertilization program is continued with limited fishing pressure, overcrowding may have disastrous consequences and result in a fish kill.

Table 4. How angler catch relates to fish population balance.	
Catch	Population Condition
- Bluegill six inches and larger - Bass average two pounds, but other sizes are caught	Populations are balanced
- Bluegill three to five inches - Bass are hard to catch	Bream population is overcrowded
- Bluegill larger than 1/3 pound - Bass less than one pound	Bass population is overcrowded
- Small crappie, sunfish, bullheads, carp, suckers, or shiners	Other species are competing with the bass and bream populations

Supplement the bream forage to get a few extra pounds on trophy bass. Managing for large bass allows intermediate and large bream to become prey to bass. A bass in the 10 pound range can easily eat a 1/3 pound bream. Bream reproduction in trophy bass ponds may diminish to a point that supplemental forage stocking is needed. Threadfin shad or fathead minnows are the best forage options for stocking Georgia ponds. These fish should be stocked in the cooler months so that they may have an opportunity to spawn in the spring. Successful spawning can occur by stocking 600 threadfin shad or 2,000 fathead minnows per acre. When a decline in forage numbers is noticed, restock as needed.

Structure in a pond helps anglers harvest more fish. While structure will not

increase the number of fish, it concentrates the ones that are there. If a pond is constructed with brush piles, ridges, or rock piles, locate the structure at different depths. After a pond has been filled, add structure such as Christmas trees, hardwood limbs, or man-made structures.

Fish populations should be monitored by angler catch records, electro-fishing, or seining. Most pond managers should be able to keep records of the size, number and species caught from their ponds (Table 4). Use the harvest goals that were set when the pond was stocked, fish the pond accordingly. All anglers who fish the pond should leave a record of the fish they catch (Figure 1). Some ponds have a mail box for depositing fishing records for the manager to check. However, when angler catch records are not kept, electro-fishing or seining can be use to check population balance. Electro-fishing requires a specially rigged boat that can go where the fish are to temporarily stun then causing them to float to the surface where they can be counted, weighed, and identified. Large mouth bass may be sampled more effectively if electro-fishing is done at night. Seining with 25 and 50 ft seines allows the shoreline fish to be sampled. Small fish can be caught in this manner to determine the success of fish spawning. Large fish are usually not caught by shore seining.

Figure 1. An example of an angler diary for keeping fishing records.

Date	Initial	Number Fishing	Time Fishing		Bass		Bluegill		Red ear	
			Start	End	No.	Wt.	No.	Wt.	No.	Wt.
6/25	GB	1	6 AM	10 AM	R 2	2 lb	K 10	1 lb	---	---

K = Number of fish kept.
R = Number of fish released.

The weight of a fish relative to a standard weight shows if the fish is getting enough to eat or not. The condition of a fish can indicate the balance between that fish and other fish in the pond of its own species and predator or prey species. For example, if a bass is thin there may be too many bass and too few bream. Use the tables 5, 6 and 7 to compare standard weights to the weight of fish caught from a pond. If the weight is within 90 percent of the standard weight, conditions are

okay. If the fish are 80 percent of less of the standard weight, improvements should be made to correct the pond management system.

Table 5. Weights of bass at different lengths for determining fish condition.

Standard weights for northern largemouth bass.	
Length, in	Weight, oz (lb)
- 12	- 14.4 (0.9)
- 14	- 20.8 (1.5)
- 16	- 35.2 (2.2)
- 18	- 51.2 (3.2)
- 20	- 72 (4.5)
- 22	- 99.2 (6.2)
- 24	- 129.6 (8.1)

Table 6. Weights of bluegill at different lengths for determining fish condition.

Standard weights for bluegill sunfish.	
Length, in	Weight, oz (lb)
- 3 (first year)	- 1.25 (0.075)
- 4.5 (second year)	- 2.5 (0.15)
- 6 (third year)	- 3.2 (0.2)
- 7.5	- 4.9 (0.3)

Table 7. Weights of crappie at different lengths for determining fish condition.

Standard weights for black crappie.	
Length, in	Weight, oz (lb)
- 5.3	- 1.2 (0.07)
- 6.3	- 1.7 (0.11)
- 8.0	- 3.5 (0.22)
- 10.0	- 7.2 (0.45)
- 11.0	- 9.2 (0.57)
- 12.0	- 12.3 (0.77)

Renovating Fish Populations

When pond fish populations are out of balance and there is no acceptable method to restore balance, the pond may need to be renovated by draining, drying, and refilling before stocking a new fish population. If the pond can be drained, some of the fish may be salvaged for future use or sale. However, most recreational fish ponds were not constructed so that fish could be seined or netted from shallow water. Ponds may have shallow puddles to holes with three or more feet of water that cannot be easily drained. In those cases, a chemical fish toxicant may be used to complete the renovation.

Rotenone can be used to eliminate unwanted fish to make a new start on balancing the pond fish population. Rotenone is effective in ponds with deep water. Several formulations containing rotenone have been developed that have different effective rates that depend on the water depth, water temperature, and fish species. Rotenone biodegrades in four days to five weeks, depending on water temperature. Avoid applications where overflow of the treated pond may drain into other fish-bearing water. Detoxify rotenone by applying an equal amount of potassium permanganate or chlorine. Rotenone products must be used according to their label instructions and are restricted use pesticides. A certified pesticide applicator must be directly involved in application of rotenone and appropriate permits and permissions must be obtained.

References:

- Anderson, R. O. and Neumann, R. M. 1996. Length, weight, and associated structural indices. Pp. 447-482. In B. R. Murphy and D.W. Willis (eds.) *Fisheries Techniques*. American Fisheries Society, Bethesda, Maryland.
- Carlander, K. D. 1977. *Handbook of freshwater fishery biology, volume two: Life history data on Centrarchid fishes of the United States and Canada*. Iowa State University Press, Ames, Iowa. 431 pp.
- Boyd, C. E. and Sowles, J. W. 1978. Nitrogen fertilization of ponds. *Transactions of the American Fisheries Society* 107(5): 737-741.
- Brunson, M. W., Stone, N., and Hargreaves, J. 1999. *Fertilization of fish ponds*. Southern Regional Aquaculture Center Publication 471. Mississippi State University, Mississippi State, Mississippi. 4 pp.

Neuman, M. 2003. Sportfish management in Alabama ponds. Alabama Department of Conservation and Natural Resources, Division of Wildlife and Fisheries. Montgomery, Alabama. 24 pp.

Prentice, J. A. and Schleckte, J. W. 2000. Performance comparison between coppernose and native Texas bluegill populations. Proceedings of the Fifty-fourth Annual Conference, Southeastern Association of Fish and Wildlife Agencies. 54:196-206.

Rice, J. S., Noble, R., and Curry, R. 2010. Pond Management Guide. AG-424 (revised). North Carolina Wildlife Resources Commission and North Carolina Cooperative Extension Service. Raleigh, North Carolina. 31 pp.

Swingle, H. S. 1950. Relationships and dynamics of balanced and unbalanced fish populations. Agricultural Experiment Station Bulletin 274. Alabama Polytechnic Institute. Auburn, Alabama. 74 pp.

Swingle, H. S. and Shell, E. W. 1971. Tables for computing relative conditions of some common freshwater fishes. Agricultural Experiment Station Circular 183. Auburn University, Alabama. 55 pp.

Wright, R. and Masser, M. 2004. Management of recreational fish ponds in Alabama. ANR 577. Alabama Cooperative Extension Service System, Alabama A&M and Auburn University. Auburn University, Alabama. 22 pp.

The University of Georgia and Ft. Valley State University, the U.S. Department of Agriculture and counties of the state cooperating. The Cooperative Extension Service offers educational programs, assistance and materials to all people without regard to race, color, national origin, age, sex or disability.

An Equal Opportunity Employer/Affirmative Action organization Committed to a Diverse Work Force

Issued in furtherance of Cooperative Extension work, Acts of May 8 and Jun 30, 1914, The University of Georgia College of Agricultural and Environmental Sciences and the U.S. Department of Agriculture cooperating.