Elmer W. Gray, Entomologist

Mosquito control operators are required to conform to the Clean Water Act (CWA) and be permitted by Georgia's National Pollutant Discharge Elimination System (NPDES). The state NPDES is managed through the Georgia Environmental Protection Division (404) 657-5947. A Notice of Intent (NOI) must be submitted to the EPD within 30-days of making a pesticide application that contacts "waters of the state." In addition, the Georgia Department of Agriculture now has a Mosquito

Control category (41). License holders in category 31 who make mosquito control treatments are now required to be licensed in Category 41. The Georgia Mosquito Control Association website (gamosquito.org) has comprehensive information about these new regulations. The University of Georgia Cooperative Extension service can be reached at (706) 542-1184 for additional information related to these matters.

Table 1. Label Recommendations of Insecticides for Use as Ultra Low Volume and Low Volume Sprays in Ground Applications for the Control of Adult Mosquitoes

		APPLIC	ATION		
INSECTICIDE/ TRADE NAME	MOA	FLOW RATE FL OZ/MIN	SPEED MPH	REMARKS AND PRECAUTIONS	
chlorpyrifos Mosquitomist One ULV	1B	3.9-7.8	5-20	Non-corrosive, quick knockdown, low odor, and low mammalian toxicity. Breaks down quickly in the environment. Good option for resistance management purposes.	
Mosquitomist Two ULV]	1.0-8.6	5-20		
Mosquitomist 1.5	1	1.4-10.5	5-20		
Pyrifos One Mist ULV EC]	1.9-15.5	5-20		
PyroFos 1.5 ULV]	1.4-10.5	5-20		
chlorpyrifos + permethrin ULV Mosquito Master 412	1B + 3	2.3-12.9	5-15		
deltamethrin DeltaGard	3	1.0-60.9	5-20	No piperonyl butoxide (pbo) in formulation, Type II pyrethroid.	
etofenprox Aqua Zenivex E20	3A	0.9-40.4	10-20	Can be diluted with water.	
Zenivex E4 RTU]	2.25-18.0	5-15	Newer product that may be helpful if resistance is a concern.	
Zenivex E20]	0.9-52.8	10-20		
malathion Fyfanon ULV	1B	1-8.6	5-20	Long time standard and still effective and cost efficient. Odor present, excites mosquitoes prior to mortality.	
naled Dibrom Concentrate	1B	0.6-1.8	5-15	Restricted Use Pesticide due to eye and skin hazard. Best choice for salt-marsh mosquitoes, typically applied aerially, rapidly degrades in the environment,	
Trumpet EC		0.8-2.2	5-15	however highly corrosive to all equipment.	
permethrin/pbo Aqualuer 20-20	3	1.6-6.2	10-20	Resistance is a concern. Areas that have used <i>permethrin</i> over several seasons should evaluate pest population for susceptibility.	
Aqua Perm-X UL 30-30	1	0.5-21.5	10-20	Excellent choice for ground applications. Variety of applications.	
Aqua-Reslin	1	0.7-35.4	10-20	Widely used and highly effective. Maintain the rate in the 0.0035 lb/A range or higher.	
Biomist 1.5 - 7.5	1	6-54	5-15		
Biomist 3 + 15	1	1.5-27	5-15		
Biomist 4 + 4		4.5-18	5-15		
Biomist 30 + 30]	1-6	5-15		
Biomist 31 + 66		1-6	5-15		

Table 1. Label Recommendations of Insecticides for Use as Ultra Low Volume and Low Volume Sprays in Ground Applications for the Control of Adult Mosquitoes

	MOA	APPLICATION			
INSECTICIDE/ TRADE NAME		FLOW RATE FL OZ/MIN	SPEED MPH	REMARKS AND PRECAUTIONS	
permethrin/pbo (continued) Envion	3	0.5-25.8	10-20	Can be diluted with water or oil.	
Envion RTU		2.4-16.5	10	Pre-mixed, water diluted, ready to use formulation.	
Evolver 4-4 ULV		4.7-75.2		Depending on dilution rate and speed.	
Evolver 30-30 ULV		1.1-2.2	10		
Aqua-Kontrol Concentrate		2-15.5	5-20	May be diluted with water.	
Kontrol 2-2		2-15.5	5-20		
Kontrol 4-4		4-16	5-20		
Kontrol 30-30		0.5-2.2	5-20		
Permanone - Ready-to-use		6-18	5-15		
Permanone 30-30		0.5-2	5-20		
Permanone 31-66		0.5-15	5-20		
Perm-X UL 4-4		1.4-105	5-20		
Perm-X UL 30-30		0.3-38.2	5-20		
Perm-X UL 31-66		0.2-33.6	5-20		
sumithrin/pbo Anvil 2 + 2 ULV	3	3.1-37.2	5-20	Good for truck applications.	
Anvil 10 + 10		0.6-7.6	5-20		
pralletron and sumithrin Duet	3	2.5-7.5	5-20	Combines two active ingredients, causes benign agitation, useful against Asian Tiger Mosquito.	

Table 2. Label recommendations of Insecticides for Use as Thermal Aerosols in Ground Applications for the Control of Adult Mosquitoes

		APPLICATION			
INSECTICIDE/ TRADE NAME			SPEED MPH	REMARKS AND PRECAUTIONS	
chlorpyrifos Mosquitomist One ULV Mosquitomist 1.5 ULV PyroFos 1.5 ULV	1B 	9.6 6.7 6.4	5	Non-corrosive, quick knockdown, low odor, and low mammalian toxicity. Breaks down quickly in the environment. Viable option for resistance management purposes.	
etofenprox Aqua Zenivex E20 Zenivex E4 RTU	3A	0.9-40.4 0.3-48.0	10-20 5-15	Apply at the rate of 0.00175-0.0070 lb ai/A.	
Zenivex E20		0.9-52.8	10-20		
malathion Fyfanon ULV	1B	1.6-2.1	5	Long time standard and still effective and cost efficient. Odor present, excites mosquitoes prior to mortality.	
naled Dibrom Concentrate	1B	1.6-2.1	5-15	Restricted Use Pesticide due to eye and skin hazard. Best choice for salt-marsh mosquitoes, typically applied aerially, rapidly degrades in the environment, however highly corrosive to all equipment.	
permethrin/pbo Aqualuer 20-20 Aqua Perm-X UL 30-30	3	0.7-25.5 0.5-21.5	10-20 10-20	Rate not to exceed 0.007 lb/A.	
Envion		2.2-25.8	10-20	Do not exceed 0.007 lb ai/A.	
Envion RTU		1.2- 32.9	5-20	Droplet size will range from 0.1-1 microns.	
Evolver 4-4 ULV			5-20	Apply in 0.0035-0.007 lb ai/A range.	
Evolver 30-30 ULV				Rates vary according to dilution, do not exceed 0.007 lb/A.	
Kontrol 2-2		10.5-32	5-20	Widely used and highly effective. Maintain dose in the 0.0035 lb/A range.	
Kontrol 4-4		21.0-64	5-20	1	
Kontrol 30-30		21.0-64	5-20		
Permanone 30-30		0.41-3	5-20	Flow rate depends on concentration. Rate not to exceed 0.007 lb permethrin/A.	
Perm-X UL 4-4		1.4-105	5-20		
Perm-X UL 30-30		0.3-38.2	5-20		
Perm-X UL 31-66		0.2-33.6	5-20		
pyrethrin/pbo Pyrenone 25-5	3	12-30	Variable	Rate not to exceed 0.0025 lb of <i>pyrethrins</i> /A. No water set back, many indoor uses.	
sumithrin/phenothrin/pbo Anvil 2+2 ULV	3, 3A	3.1-37.2	5-20	Good for truck applications.	
Anvil 10+10 ULV		0.6-7.6	5-20		

Table 3. Label Recommendations of Insecticides for Use in Aerial Applications for Control of Adult Mosquitoes.

INSECTICIDE/ TRADE NAME	MOA	GROSS AMOUNT PER ACRE (OZ)	REMARKS AND PRECAUTIONS
<i>chlorpyrifos</i> Mosquitomist One ULV	1B	1.28	Non-corrosive, quick knockdown, low odor, and low mammalian toxicity. Breaks down quickly in the environment.
PyroFos 1.5 ULV	1	0.87	Viable option for resistance management purposes.
etofenprox Aqua Zenivex E20	3A		Apply diluted or undiluted at rates up to 0.007 lb ai/A. Minimum application height 100 ft.
Zenivex E4 RTU			
Zenivex E20			Apply at 100-300 ft altitude.
<i>malathion</i> Fyfanon ULV	1B	1-3	Long time standard and still effective and cost efficient. Odor present, excites mosquitoes prior to mortality. New lower rate recently approved by EPA.
naled Dibrom Concentrate	1B	0.5-1	Restricted Use Pesticide due to eye and skin hazard. Best choice for salt-marsh mosquitoes, typically applied aerially, rapidly degrades in the environment, however, highly corrosive to all equipment.
Trumpet EC		0.6-1.2	consider to an equipment.
permethrin/pbo AquaHalt Aqualuer 20-20	3	0.28-0.76	Water based, labeled for use over crops. Apply diluted or undiluted at rates up to 0.007 lb ai/A.
Aqua-Reslin	1	0.25-0.5	Widely used and highly effective. Maintain dose in the 0.0035 lb/A range.
Biomist 1.5 + 7.5 ULV	1	9	
Biomist 4.0 + 4.0 ULV	-	3	··· ·
Biomist 12 + 60 ULV	1	1-3	
Biomist 30 + 30 ULV	.1	0.33-1	
Biomist 31 + 66 ULV]	0.33-1	
Envion]		Apply diluted or undiluted at rates up to 0.007 lb permethrin/pbo/A.
Envion RTU		up to 2.72	Apply at altitudes between 100-300 ft.
Evolver 4-4 ULV		3.1	Do not exceed 3.1 fl oz/A.
Evolver 30-30 ULV		0.05-0.36	
Aqua-Kontrol Concentrate		up to 1.5	Apply at sufficient air speed to deliver the appropriate amount of a.i./A (from 0.00175-0.007 lb).
Kontrol 4 + 4		up to 3	
Kontrol 30 + 30		up to 0.35	
Permanone 31-66		0.16-0.6	
Perm-X UL 4-4		up to 3.1	
Perm-X UL 30-30		up to 0.32	
Perm-X UL 31-66	1 2	up to 0.36	Annihom to 0.0025 lb of condition Annihom to design to the state of th
pyrethrins/pbo Pyrenone 25-5	3		Apply up to 0.0025 lb of <i>pyrethrins</i> /A with equipment designed and operated to produce a ULV spray application.
sumithrin/pbo Anvil 2 + 2 ULV Anvil 10 + 10 ULV	3	0.21-0.62	
prallethrin and sumithrin Duet	3	0.41-1.24	Combines two active ingredients, causes benign agitation, option for Asian Tiger Mosquito.

Table 4. Label Recommendations of Insecticides for Use in Ground Applications for Larval Mosquito Control.

INSECTICIDE/ TRADE NAME	MOA	GROSS AMOUNT PER ACRE (OZ)	REMARKS AND PRECAUTIONS
mineral oil BVA2 Mosquito Larvicide Oil		1-5 gal	Surface film material. Effective against both larvae and pupae, low labor input required. Use higher rates for areas with more vegetation or pollution.
Cocobear Mosquito Larvicide Oil		3-5 gal	
larvicide oil Kontrol Mosquito Larvicide		1-5 gal	
methoprene Altosid Liquid Larvicide	7A	3-4 fl oz	Significant residual effect possible, up-front cost more, can save labor costs. Approved for use in stock waters. Newly approved for ULV applications. Used to prepare Altosand granular formulation.
Altosid Liquid Larvicide Concentrate		0.75-1 fl oz	
Altosid 30-Day Briquets		1/100 sq ft	
Altosid XR Briquets		1 briquet/100-200 sq ft	For areas up to 2 ft in depth; use additional briquette for areas over 2 ft in depth. Effective up to 150 days.
Altosid P35		2.5-20 lb	New product, spherical design to improve calibration and treatment accuracy, 35 days of residual control.
Altosid Pellets		2.5-10 lb	Rate depends on species targeted. Use lower rates when water is shallow or vegetation, and/or pollution is minimal. Effective up to 30 days.
Altosid Pellets Water Soluble Pouch (WSP)		Up to 135 sq ft	
Altosid SBG II		3-20 lb	Rate depends on target species.
Altosid XR-G		5-20 lb	May be applied prior to flooding.
MetaLarv S-PT		2.5-10 lb	Use lower rates when water is shallow, vegetation, and/or pollution are minimal and mosquito populations are low.
Bacillus thuringiensis subsp. israelensis Aquabac XT 12AS	11A	1-2 pt	Biological control agent, very effective, environmentally sound, store products in cool areas, and use by expiration date. Timing of application critical for maximum effectiveness. Use higher label rates when large
Aquabac 200G		2.5-10 lb	larval populations are present, larvae are in late 3rd or 4th instars or local experience has demonstrated that
Aquabac 400G		1.25-8 lb	higher rates are required to achieve high levels of mortality.
Bactimos/Summit Briquets/Dunks		1 briquet/100 sq ft	
Teknar HP-D		0.25-1 pt	
Vectobac CG		2.5-10 lb	
Vectobac 12AS		5-100 gal	
Vectobac G		2-20 lb	
Fourstar Multi-Brood Controlled Release Granule		7.5-20 lb	
Fourstar Bti Briquet 150-Day		1 briquet/100 sq ft	Alternate wetting and drying will not reduce effectiveness, performs optimally in shaded areas.
Bacillus sphaericus Vectolex WSP	11B	1 pouch/50 sq ft	Biological control agent with residual effect. Targets specific species, especially Culex. Not recommended for salt-marsh mosquitoes.
Vectolex CG		5-20 lb	
(Waste Tires)		20-80 lb or 0.5-2 lb/1000 sq ft	
Vectolex WDG		0.5-1.5 lb	

Table 4. Label Recommendations of Insecticides for Use in Ground Applications for Larval Mosquito Control.

INSECTICIDE/ TRADE NAME	MOA	GROSS AMOUNT PER ACRE (OZ)	REMARKS AND PRECAUTIONS
Bacillus thiringiensis subsp. israelensis and Bacillus sphaericus	11A	5-20 lb/A	Effective against all pest species of mosquitoes, will reduce likelihood of resistance.
Vectomax FG	+	20-80 lb/A for tire piles	Approved for use in organic crops and sensitive areas.
Vectomax WSP	11B	1 pouch/50 sq ft	
Vectomax G		5-20 lb/A 20-80 lb/A for tire	
		piles	
Fourstar 45-Day Briquet		1 briquet/100 sq ft	
Fourstar 90-Day Briquet		1 briquet/100 sq ft	
Fourstar 180-Day Briquet		1 briquet/100 sq ft	
Fourstar Multi-Brood Granule		7.5-20 lb/A	Effective up to 4 weeks.
Fourstar Multi-Brood Controlled Release Granule		5-20 lb/A	
Fourstar Single Brood Bti Sand Granule		3-10 lb	
Fourstar Water Soluble Pouch		1 pouch/50 sq ft of larval habitat	
Bacillus thuringiensis subsp. israelensis and (s)-methoprene	11A	1.25-10 lb/A	Post-flood – Use higher rates when water is heavily polluted.
Vectoprime	+ 7A	10-20 lb/A	Pre-flood – 7 days pre-flood use 10-20 lb/A; and for 7-14 days pre-flood. Wider application window for early or late instars.
Duplex – G		2.5-20 lb/A	Heavy, uniform granule, improves calibration and penetration into habitats.
spinosad	5		
Natular G – Single Brood Granule		3.5-9 lb/A	Registered as a reduced risk pesticide by EPA.
Natular G-30		5-20	Multiple-brood extended release granule.
Natular 2EC – Single Brood Liquid		1.3-2.8 fl oz/A	
Natular T30		1 tablet/100 sq ft	Multiple-brood, 30-day tablet.
Natular XRT		1 tablet/100 sq ft	Multiple-brood extended release tablet.

Table 5. Label Recommendations of Insecticides for Use in Aerial Applications for Control of Larval Mosquitoes.

INSECTICIDE/ TRADE NAME	MOA	GROSS AMOUNT PER ACRE (OZ)	REMARKS AND PRECAUTIONS
mineral oil BVA 2 Mosquito Larvicide Oil		3-5 gal	Surface film material. Not for use in environmentally sensitive areas. Surface film product that controls larvae and pupae.
Cocobear Mosquito Larvicide Oil Kontrol Mosquito Larvicide Oil		3-5 gal Aerial: 2-4 gal Ground: 1-5 gal	Rate depends on water surface conditions and vegetative density.

Table 5. Label Recommendations of Insecticides for Use in Aerial Applications for Control of Larval Mosquitoes.

INSECTICIDE/ TRADE NAME	MOA	GROSS AMOUNT PER ACRE (OZ)	REMARKS AND PRECAUTIONS
methoprene	7A		Effective against both larvae and pupae, low labor input required.
Altosid Liquid Larvicide		3-4 fl oz	
Altoside Liquid Larvicide Concentrate		0.75-1 fl oz	
Altosid P35		2.5-20 lb	New product, spherical design to improve calibration and treatment accuracy, 35 days of residual control.
Altosid pellets		2.5-10 lb	
Altosid XR-G	Ì	1/200 sq ft	
Altosand G	Ì	8-10 lb	
MetaLarv S-PT		2.5-10 lb	Use lower rates when water is shallow, vegetation and/or pollution are minimal and mosquito populations are low.
Bacillus thuringiensis subsp. israelensis	11A		Biological control agent, very effective, environmentally sound, store products in cool areas and use by expiration
Aquabac XT		0.25-2 pt	date. Timing of application critical for maximum effectiveness. Granular formulations can be applied pre- or post-hatch.
Summit B.t.i. Granules		2.5-20 lb	nacii.
Teknar HPD		0.25-1 pt	
Vectobac 12 AS		0.25-2 pt	
Vectobac CG		2.5-10 lb	
Vectobac G		2.5-5 lb	
Vectobac WDG		1.75-14 oz	
Fourstar Single Brood Bti Sand Granule		3-20 lb	
Fourstar Multi-brood Controlled Release Granule		7.5-20 lb	
Sustain MBG		5-20 lb	
Bacillus sphaericus Vectolex CG	11B	5-20 lb	Biological control agent with residual effect. Targets specific species, especially Culex.
Vectolex WDG	ĺ	0.5-1.5 lb	
Vectolex WSP		1 pouch/50 sq ft	Water soluble pouch, good choice for catch basins.
Bacillus thiringiensis subsp. israensis and	11A		Effective against all pest species of mosquitoes, will reduce likelihood of resistance.
Bacillus sphaericus	+ 11B	5-20 lb	
Vectomax CG	ПВ	20-80 lb for tire piles	
Vectomax G		5-20 lb	
	ļ	20-80 lb for tire piles	
Duplex – G		2.5-20 lb	
Fourstar Multi-Brood Granule		5-20 lb	Effective up to 40 days.
Fourstar Multi-Brood Controlled Release Granule		5-20 lb	
spinosad Natular G	5	3.5-9 lb	Registered as a reduced risk pesticide by EPA.
Natular XRG		5-20 lb	
Natular 2EC	,	1.3-2.8 fl oz/A	

Table 6. Toxicity of Insecticides Used in Mosquito Control Compared to Some Other Common Insecticides and Chemicals.

		ACUTE ORAL LD ₅₀ IN MG TECH/KG FOR RATS BY REFERENCE NUMBER					
COMMON NAME	TRADE NAME	1*	2**	3***			
allethrin	b	310	680-1,000	920			
aspirin		1,900ª					
bti	b	non-toxic		non-toxic			
chlorpyrifos	Dursban	135	82-245	163/135			
dichlorvos	DDVP	56	25-170				
fenitrothion	Sumithion	490	250-740	300/800			
fenthion	Baytex	190	255-740	375/290			
malathion	Fyfanon	1,375	885-2,800	1,375/1,000			
methoprene	b	34,600	> 34,600	> 34,600			
methoxychlor	b	6,000	5,000-7,000	> 6,000			
monomolecular firm	Agnique	20,000		> 20,000			
naled	Dibrom	430	250-430	430			
phenothrin	Sumithrin	10,000	> 10,000	> 10,000			
piperonyl butoxide			6,150-7,500	11,500			
pyrethrin	b	200	200-2,600	820-1,870			
resmethrin	SBP-1382	2,500	1,500	4,240			
salt		4,500a					
sugar		27,000ª					
temephos	b	2,030	1,000-13,000	8,600-1,300			

^a The Bottom Line. Dow Chemical Co., Midland, MI. January 1981, 7 pp. (from: Hodge, H.C. and W.C. Downs, Toxicol. Appl. Pharmacol. 3:689-695, 1961).

^b Manufactured under several trade names.

^{*} Thompson, W.T. 1985. Agricultural chemicals, Book 1 Insecticides, 1985-86 Revision. Thompson Pub., Fresno, CA 255 pp.

^{**} Kenaga, E.E., and R.W. Morgan. 1978. Commercial and Experimental Organic Insecticides. 1978 Revision, Entomol. Soc. of Am. Pub. 78-1. 79 pp.

^{***} From manufacturers technical data.