

PEANUT: PEANUT INSECT CONTROL

Mark R. Abney, Research and Extension Entomologist

PEST	INSECTICIDE	MOA	AMOUNT OF FORMULATION PER ACRE	LB ACTIVE INGREDIENT PER ACRE	REI/PHI (Hours or Days)	REMARKS
Beet Armyworm ¹	<i>chlorantraniliprole</i> Prevathon	28	14-20 fl oz	0.047-0.067	4 H/ 1 D	Do not apply more than 0.2 lb ai of <i>chlorantraniliprole</i> /A/ year.
	<i>diflubenzuron</i> Dimilin 2L	15	4-8 fl oz	0.06-0.125	12 H/ 28 D	Do not make more than 3 applications/season.
	<i>indoxacarb</i> Steward 1.25EC	22	9.2-11.3 fl oz	0.09-0.11	12 H/ 14 D	Do not apply more than 45 fl oz/A/season. Minimum interval between treatments is 5 days.
	<i>methoxyfenozide</i> Intrepid 2F	18	6-10 fl oz	0.09-0.16	4 H/ 7 D	
	<i>novaluron</i> Diamond 0.83 EC	15	6-12 fl oz	0.039-0.077	12 H/ 28 D	Do not apply more than 36 fl oz/A/season. Do not feed treated peanut hay or vines to livestock.
	<i>spinetoram</i> Radiant 1SC	5	3-8 fl oz	0.023-0.063	4 H/ 3 D	Do not apply more than a total of 24 fl oz of Radiant SC (0.188 lb ai <i>spinetoram</i>)/A/year. Do not allow grazing of peanut hay.
	<i>spinosad</i> Blackhawk 36WG	5	1.7-3.3 oz	0.038-0.074	4 H/ Harvest 3 D Forage 14 D	Do not apply more than 12.4 oz/A/crop. Do not make applications less than 7 days apart.
Burrower Bug ⁵	<i>chlorpyrifos</i> Chlorpyrifos 15G	1B	13.6 lb	2	24 H/ 21 D	Maximum single application rate is 15 oz/1000 ft of row. Do not exceed 26.4 lb Chlorpyrifos 15G/A/season. For banded applications use a 10-18" band.
Corn Earworm	<i>chlorantraniliprole</i> Prevathon	28	14-20 fl oz	0.047-0.067	4 H/ 1 D	Do not apply more than 0.2 lb ai of <i>chlorantraniliprole</i> /A/year.
	<i>indoxacarb</i> Steward 1.25EC	22	9.2-11.3 fl oz	0.09-0.11	12 H/ 14 D	Do not apply more than 45 fl oz/A/season. Minimum interval between treatments is 5 days.
	<i>methomyl</i> Lannate 90SP Lannate 2.4LV	1A	0.25-1 lb 0.75-3 pt	0.225-0.9	48 H/ 21 D	Do not make more than 3 applications. Do not feed treated vines to livestock.
	<i>spinetoram</i> Radiant 1SC	5	3-8 fl oz	0.023-0.063	4 H/ 3 D	Do not apply more than a total of 24 fl oz of Radiant SC (0.188 lb ai <i>spinetoram</i>)/A/year. Do not allow grazing of peanut hay.
	<i>pyrethroids</i> <i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	1.8-2.4 fl oz	0.014-0.019	12 H/ 14 D	Do not exceed 3 applications/season of 2.8 fl oz/10-day interval.
	<i>bifenthrin</i> Brigade 2EC		2.1-6.4 fl oz	0.033-0.1	12 H/ 14 D	Do not apply more than 0.5 lb ai/A/season. Do not feed peanut hay to livestock.
	<i>esfenvalerate</i> Asana XL 0.66EC		2.9-5.8 fl oz	0.015-0.03	12 H/ 21 D	Do not exceed 0.15 lb ai/A/season. Do not feed or graze livestock on treated vines.
	<i>lambda-cyhalothrin</i> Warrior II Zeon 2.08CS		1.28-1.92 fl oz	0.02-0.03	24 H/ 14 D	Do not apply more than 1 pt/A/season. Do not graze livestock in treated areas or use treated vines for animal feed.
	<i>zeta-cypermethrin</i> Mustang Maxx 0.8EC		3.2-4 oz	0.02-0.025	12 H/ 7 D	Do not apply more than 0.15 lb ai/season. Do not graze livestock in treated areas. Do not use treated vines for animal feed.

PEANUT INSECT CONTROL

PEST	INSECTICIDE	MOA	AMOUNT OF FORMULATION PER ACRE	LB ACTIVE INGREDIENT PER ACRE	REI/PHI (Hours or Days)	REMARKS
Cutworm ¹	<i>indoxacarb</i> Steward 1.25EC	22	9.2-11.3 fl oz	0.09-0.11	12 H/ 14 D	Do not apply more than 45 fl oz/A/season. Minimum interval between treatments is 5 days.
	<i>methomyl</i> Lannate 90SP Lannate 2.4LV	1A	0.5-1 lb 1.5-3 pt	0.45-0.9	48 H/ 21 D	Spray late in the afternoon for maximum efficacy. Do not feed treated vines to livestock.
	<i>pyrethroids</i> <i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	1-1.8 fl oz	0.008-0.014	12 H/ 14 D	Do not exceed 3 applications per season of 2.8 fl oz/10-day intervals.
	<i>bifenthrin</i> Brigade 2EC		2.1-6.4 fl oz	0.033-0.1	12 H/ 14 D	Do not apply more than 0.5 lb ai/A/season. Do not feed peanut hay to livestock.
	<i>esfenvalerate</i> Asana XL 0.66EC		9.6 oz	0.05	12 H/ 21 D	Do not exceed 0.15 lb ai/A/season. Do not feed or graze livestock on treated vines.
	<i>lambda-cyhalothrin</i> Warrior II Zeon 2.08CS		1.28-1.92 fl oz	0.015-0.025	24 H/ 14 D	Do not apply more than 1 pt/A/season. Do not graze livestock in treated areas or use treated vines for animal feed.
	<i>zeta-cypermethrin</i> Mustang Maxx 0.8EC		1.28-4 oz	0.008-0.025	12 H/ 7 D	Do not apply more than 0.15 lb ai/season. Do not graze livestock in treated areas. Do not use treated vines for animal feed.
Fall Armyworm ¹	<i>chlorantraniliprole</i> Prevathon	28	14-20 fl oz	0.047-0.067	4 H/ 1 day	Do not apply more than 0.2 lb ai of <i>chlorantraniliprole</i> /A/year.
	<i>diflubenzuron</i> Dimilin 2L	15	4-8 fl oz	0.06-0.125	12 H/ 28 D	Do not make more than 3 applications/season.
	<i>indoxacarb</i> Steward 1.25EC	22	9.2-11.3 fl oz	0.09-0.11	12 H/ 14 D	Do not apply more than 45 fl oz/A/season. Minimum interval between treatments is 5 days.
	<i>methomyl</i> Lannate 90SP Lannate 2.4LV	1A	0.25-0.5 lb 0.75-1.5 pt	0.225-0.45	48 H/ 21 D	Do not feed treated vines to livestock.
	<i>methoxyfenozone</i> Intrepid 2F	18	6-10 fl oz	0.09-0.16	4 H/ 14 D	Do not make more than 3 applications/year.
	<i>novaluron</i> Diamond 0.83 EC	15	6-12 fl oz	0.039-0.077	12 H/ 28 D	Do not apply more than 36 fl oz/A/season. Do not feed treated peanut hay or vines to livestock.
	<i>spinetoram</i> Radiant 1SC	5	3-8 fl oz	0.023-0.063	4 H/ 3 D	Do not apply more than a total of 24 fl oz of Radiant SC (0.188 lb ai <i>spinetoram</i>)/A/year. Do not allow grazing of peanut hay.
	<i>spinosad</i> Blackhawk 36WG	5	1.7-3.3 oz	0.038-0.074	4 H/ Harvest 3 D Forage 14 D	Do not apply more than 12.4 oz/A/crop. Do not make applications less than 7 days apart.
Green Cloverworm ¹	<i>methoxyfenozone</i> Intrepid 2F	18	6-10 fl oz	0.09-0.16	4 H/ 14 D	Do not make more than 3 applications/year.
	<i>novaluron</i> Diamond 0.83 EC	15	6-12 fl oz	0.039-0.077	12 H/ 28 D	Do not apply more than 36 fl oz/A/season. Do not feed treated peanut hay or vines to livestock.

PEANUT INSECT CONTROL

PEST	INSECTICIDE	MOA	AMOUNT OF FORMULATION PER ACRE	LB ACTIVE INGREDIENT PER ACRE	REI/PHI (Hours or Days)	REMARKS
Green Cloverworm ¹ (continued)	<i>spinetoram</i> Radiant 1SC	5	3-8 fl oz	0.023-0.063	4 H/ 3 D	Do not apply more than a total of 24 fl oz of Radiant SC (0.188 lb ai <i>spinetoram</i>)/A/year. Do not allow grazing of peanut hay.
	<i>spinosad</i> Blackhawk 36 WG	5	1.7-3.3 oz	0.038-0.074	4 hours/ Harvest 3 D Forage 14 D	Do not apply more than 12.4 oz/A/crop. Do not make applications less than 7 days apart.
Lesser cornstalk borer ²	<i>chlorantraniliprole</i> Prevathon	28	14-20 fl oz	0.047-0.067	4 H/ 1 D	Do not apply more than 0.2 lb ai of <i>chlorantraniliprole</i> /A/year.
	<i>chlorpyrifos</i> Chlorpyrifos 15G	1B	13.6 lb banded	2	24 H/ 21 D	Maximum single application rate is 15 oz/1000 ft of row. Do not exceed 26.4 lb Chlorpyrifos 15G/A/season. For rescue treatment apply in a 10-18" band over the fruiting zone.
	<i>novaluron</i> Diamond 0.83 EC	15	6-12 fl oz	0.039-0.077	12 H/ 28 D	Do not apply more than 36 fl oz/A/season. Do not feed treated peanut hay or vines to livestock.
Potato leafhopper	<i>acephate</i> Orthene 97	1B	12-16 oz	0.72-0.97	24 H/ Digging 14 D	Do not feed treated forage or hay to livestock or allow animals to graze treated areas.
	<i>carbaryl</i> Sevin XLR or 4F Sevin 80S	1A	1 qt 1.25 lb	1	12 H/ 14 D	
	<i>pyrethroids</i> <i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	1-1.8 fl oz	0.008-0.014	12 H/ 14 D	Do not exceed 3 applications per season of 2.8 fl oz/10-day intervals.
	<i>bifenthrin</i> Brigade 2EC		2.1-6.4 fl oz	0.033-0.1	12 H/ 14 D	Do not apply more than 0.5 lb ai/A/season. Do not feed peanut hay to livestock.
	<i>esfenvalerate</i> Asana XL 0.66EC		2.9-5.8 fl oz	0.015-0.03	12 H/ 14 D	Do not exceed 0.15 lb ai/A/season. Do not feed or graze livestock on treated vines.
	<i>lambda-cyhalothrin</i> Warrior II Zeon 2.08CS		0.96-1.6 fl oz	0.015-0.025	24 H/ 7 D	Do not apply more than 1 pt/A/season. Do not graze livestock in treated areas or use treated vines for animal feed.
	<i>zeta-cypermethrin</i> Mustang Maxx 0.8EC		1.76-4 oz	0.011-0.025	12 H/ -	Do not apply more than 0.15 lb ai/season. Do not graze livestock in treated areas. Do not use treated vines for animal feed.
Rednecked peanut worm	<i>spinetoram</i> Radiant 1SC	5	3-8 fl oz	0.023-0.063	4 H/ 3 D	Do not apply more than a total of 24 fl oz of Radiant SC (0.188 lb ai <i>spinetoram</i>)/A/year. Do not allow grazing of peanut hay.
	<i>pyrethroids</i> <i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	1-1.8 fl oz	0.008-0.014	12 H/ 14 D	Do not exceed 3 applications per season of 2.8 fl oz/10-day intervals.
	<i>bifenthrin</i> Brigade 2EC		2.1-6.4 fl oz	0.033-0.1	12 H/ 14 D	Do not apply more than 0.5 lb ai/A/season. Do not feed peanut hay to livestock.
	<i>esfenvalerate</i> Asana XL 0.66EC		2.9-5.8 fl oz	0.015-0.03	12 H/ 21 D	Do not exceed 0.15 lb ai/A/season. Do not feed or graze livestock on treated vines.
	<i>lambda-cyhalothrin</i> Warrior II Zeon 2.08CS		0.96-1.6 fl oz	0.015-0.025	24 H/ 14 D	Do not apply more than 1 pt/A/season. Do not graze livestock in treated areas or use treated vines for animal feed.
	<i>zeta-cypermethrin</i> Mustang Maxx 0.8EC		1.28-4 oz	0.008-0.025	12 H/ 7 D	Do not apply more than 0.15 lb ai/season. Do not graze livestock in treated areas. Do not use treated vines for animal feed.

PEANUT INSECT CONTROL

PEST	INSECTICIDE	MOA	AMOUNT OF FORMULATION PER ACRE	LB ACTIVE INGREDIENT PER ACRE	REI/PHI (Hours or Days)	REMARKS
Southern Armyworm ¹	<i>spinetoram</i> Radiant 1SC	5	3-8 fl oz	0.023-0.063	4 H/ 3 D	Do not apply more than a total of 24 fl oz of Radiant SC (0.188 lb ai <i>spinetoram</i>)/A/year. Do not allow grazing of peanut hay.
Southern corn rootworm ²	<i>chlorpyrifos</i> Chlorpyrifos15G	1B	13.6 lb banded	2	24 H/ 21 D	Maximum single application rate is 15 oz/1000 ft of row. Do not exceed 26.4 lb Chlorpyrifos15G/A/season. Do not feed peanut forage or hay to meat or dairy animals.
Soybean looper ¹	<i>chlorantraniliprole</i> Prevathon	28	20 fl oz	0.067	4 H/ 1 D	Do not apply more than 0.2 lb ai of <i>chlorantraniliprole</i> /A/year.
	<i>indoxacarb</i> Steward 1.25EC	22	9.2-11.3 fl oz	0.09-0.11	12 H/ 14 D	Do not apply more than 45 fl oz/A/season. Minimum interval between treatments is 5 days.
	<i>methoxyfenozide</i> Intrepid 2F	18	6-10 fl oz	0.09-0.16	4 H/ 7 D	Do not make more than 3 applications/year.
	<i>novaluron</i> Diamond 0.83 EC	15	6-12 fl oz	0.039-0.077	12 H/ 28 D	Do not apply more than 36 fl oz/A/season. Do not feed treated peanut hay or vines to livestock.
	<i>spinetoram</i> Radiant 1SC	5	3-8 fl oz	0.023-0.063	4 H/ 3 D	Do not apply more than a total of 24 fl oz of Radiant SC (0.188 lb ai <i>spinetoram</i>)/A/year. Do not allow grazing of peanut hay.
	<i>spinosad</i> Blackhawk 36WG	5	1.7-3.3 oz	0.038-0.074	4 H/ Harvest 3 D Forage 14 D	Do not apply more than 12.4 oz/A/crop. Do not make applications less than 7 days apart.
Spider mite	<i>fenpropathrin</i> Danitol 2.4EC	3A	10.6-16 fl oz	0.2-0.3	24 H/ Digging 14 D	Do not feed treated forage or dried hay within 14 days of last application. Do not exceed 0.8 lb ai/A /season.
	<i>propargite</i> Comite 6.55EC	12C	2 pt	1.64	48 H/ 14 D	Do not apply more than twice per season. Do not graze or feed livestock on treated areas or cut treated forage for hay. When temperatures are greater than 90°F with high humidity, some leaf phytotoxicity may occur.
	Comite II 6EC		2.25 pt	1.68	48 H/ 14 D	Do not apply more than twice per season. Use a minimum of 20 gal/A for ground application and 5 gal/A for aerial application.
	Omite 30W		3-5 lb	0.9-1.5	48 H/ 14 D	Do not apply more than twice per season. Do not graze or feed livestock on treated areas or cut treated forage for hay. Do not plant unregistered crops within 6 months of last application.
Three cornered alfalfa hopper	<i>carbaryl</i> Sevin XLR or 4F Sevin 80S	1A	1 qt 1.25 lb	1	12 H/ 14 D	
	<i>pyrethroids</i> <i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	1.8-2.4 fl oz	0.014-0.019	12 H/ 14 D	Do not exceed 3 applications per season of 2.8 fl oz/10-day intervals.
	<i>bifenthrin</i> Brigade 2EC		2.1-6.4 fl oz	0.033-0.1	12 H/ 14 D	Do not apply more than 0.5 lb ai/A/season. Do not feed peanut hay to livestock.
	<i>lambda-cyhalothrin</i> Warrior II Zeon 2.08CS		0.96-1.6 fl oz	0.015-0.025	24 H/ 14 D	Do not apply more than 1 pt/A/eason. Do not graze livestock in treated areas or use treated vines for animal feed.

PEANUT INSECT CONTROL

PEST	INSECTICIDE	MOA	AMOUNT OF FORMULATION PER ACRE	LB ACTIVE INGREDIENT PER ACRE	REI/PHI (Hours or Days)	REMARKS
Thrips ³	<i>acephate</i> Orthene 97	1B	6-12 oz	0.36-0.73	24 H/ Digging 14 D	Do not feed treated forage or hay to livestock or allow animals to graze treated areas. Good results have been obtained by tank mixing with early post-emergence herbicides such as <i>paraquat</i> combinations in lieu of in-furrow treatments. However, timing is critical for adequate thrips control and may not coincide with optimum timing for post-emergence weed control.
	<i>imidacloprid</i> Admire Pro 4.6F	4A	7-10.5 fl oz	0.25-0.377	12 H/ 14 D	Apply as an in-furrow spray at planting. Do not apply more than 10.5 oz/crop season. Do not apply to Virginia type varieties.
	<i>phorate</i> Thimet 20G ⁴	1B	5-7.5 lb	1	48 H/ See note	Apply in the furrow at planting. Do not graze or feed treated hay or forage to livestock. In furrow applications for single rows use 5.5 oz of 20G/1000 ft of row. Do not exceed 7.5 lbs/A for twin rows. Soil moisture is necessary for adequate uptake. Young seedlings may exhibit varying degrees of leaf damage.
Tobacco budworm ¹	<i>chlorantraniliprole</i> Prevathon	28	14-20 fl oz	0.047-0.067	4 H/ 1 D	Do not apply more than 0.2 lb ai of <i>chlorantraniliprole</i> /A/year.
	<i>indoxacarb</i> Steward 1.25 EC	22	9.2-11.3 fl oz	0.09-0.11	12 H/ 14 D	Do not apply more than 45 fl oz/A/season. Minimum interval between treatments is 5 days.
	<i>spinetoram</i> Radiant 1SC	5	3-8 fl oz	0.023-0.063	4 H/ 3 D	Do not apply more than a total of 24 fl oz of Radiant SC (0.188 lb ai <i>spinetoram</i>)/A/year. Do not allow grazing of peanut hay.
	<i>spinosad</i> Blackhawk 36WG	5	1.7-3.3 oz	0.038-0.074	4 H/ Harvest 3 D Forage 14 D	Do not apply more than 12.4 oz/A/crop. Do not make applications less than 7 days apart.
NOTE: Lannate as applied for corn earworm gives good control.						
Velvetbean caterpillar ¹	<i>diflubenzuron</i> Dimilin 2L	15	2-4 fl oz	0.03-0.06	12 H/ Harvest 28 D	Do not make more than 3 applications/season.
	<i>methoxyfenozide</i> Intrepid 2F	18	6-10 fl oz	0.09-0.16	4 H/ 7 D	Do not make more than 3 applications/year.
	<i>novaluron</i> Diamond 0.83 EC	15	6-12 fl oz	0.039-0.077	12 H/ 28 D	Do not apply more than 36 fl oz/A/season. Do not feed treated peanut hay or vines to livestock.
	<i>pyrethroids</i> <i>beta-cyfluthrin</i> Baythroid XL 1EC	3A	1-1.8 fl oz	0.008-0.014	12 H/ 14 D	Do not exceed 3 applications per season of 2.8 fl oz/10-day intervals.
	<i>bifenthrin</i> Brigade 2EC		2.1-6.4 fl oz	0.033-0.1	12 H/ 14 D	Do not apply more than 0.5 lb ai/A/season. Do not feed peanut hay to livestock.
	<i>esfenvalerate</i> Asana XL 0.66EC		2.9-5.8 fl oz	0.015-0.03	12 H/ 21 D	Do not exceed 0.15 lb ai/A/season. Do not feed or graze livestock on treated vines. Suspected resistance has been observed in lower SW Georgia.
	<i>lambda-cyhalothrin</i> Warrior II Zeon 2.08CS		0.96-1.6 fl oz	0.015-0.025	24 H/ 14 D	Do not apply more than 1 pt/A/season. Do not graze livestock in treated areas or use treated vines for animal feed. Suspected resistance has been observed in lower SW Georgia.

PEANUT INSECT CONTROL

PEST	INSECTICIDE	MOA	AMOUNT OF FORMULATION PER ACRE	LB ACTIVE INGREDIENT PER ACRE	REI/PHI (Hours or Days)	REMARKS
Velvetbean caterpillar ¹ (continued)	<i>zeta-cypermethrin</i> Mustang Maxx 0.8EC		1.28-4 oz	0.008-0.025	12 H/ 7 D	Do not apply more than 0.15 lb ai/season. Do not graze livestock in treated areas. Do not use treated vines for animal feed.
	<i>spinetoram</i> Radiant 1SC	5	3-8 fl oz	0.023-0.063	4 H/ 3 D	Do not apply more than a total of 24 fl oz of Radiant SC (0.188 lb ai <i>spinetoram</i>)/A/year. Do not allow grazing of peanut hay.
	<i>spinosad</i> Blackhawk 36WG	5	1.7-3.3 oz	0.038-0.074	4 H/ Harvest 3 D Forage 14 D	Do not apply more than 12.4 oz/A per crop. Do not make applications less than 7 days apart.
Wireworm ²	<i>chlorpyrifos</i> Chlorpyrifos 15G	1B	13.6 lb banded	2	24 H/ 21 D	Suppression only. Maximum single application rate is 15 oz/1000 ft of row. Do not exceed 26.46 lb Chlorpyrifos 15G/A/ season. If used in combination with Lorsban 4E do not exceed 4 lb ai/crop season. For banded applications, use a 10-18" band. Aerial application is prohibited.
	Lorsban 75WG		2.67 lb	2	24 H/ —	Suppression only. Broadcast pre-plant and incorporate.
	Lorsban 4E		4 pt	2	24 H/ —	Suppression only. Broadcast pre-plant and incorporate.

Premixed or Co-Packed Insecticide Products:

Products listed below are available as premixes or co-packages of two insecticidal active ingredients. When using premixed or co-packaged products, be sure the use of all of the active ingredients is necessary. Unnecessary applications or use of reduced rates of an active ingredient may lead to or intensify insecticide resistance.

imidacloprid, cyfluthrin (Leverage 2.8 fl oz/A)

imidacloprid, fluopyram (Velum Total 18 fl oz/A)

lambda-cyhalothrin, chlorantraniliprole (Besiege 5-10 fl oz/A)

methoxyfenozide, spinetoram (Intrepid Edge 4-8 fl oz/A)

¹ The treatment threshold for combined foliage feeders is 4-8 per foot of row depending on the size and condition of the peanut plants. Use a lower threshold for very young plants or plants that are stressed from other factors. Use a higher threshold for healthy plants with ample vine growth.

² Preventive treatments are usually more effective than rescue treatments. The need for a rescue treatment should be dependent on the presence of the insect pest, not just damage. Rainfall or irrigation is necessary after application to obtain adequate control.

³ Thrips control is recommended only during the first 3-4 weeks after emergence and is more important when herbicide-induced stress also occurs during early season growth. Attempts to control tomato spotted wilt by controlling the thrips vectors are not economically justified.

⁴ Suppresses tomato spotted wilt.

⁵ Risk of burrower bug is generally reduced in conventional tillage situations.

PEANUT DISEASE CONTROL

Bob Kemerait, Extension Plant Pathologist, Tim Brenneman and Albert Culbreath, Plant Pathologists

PEST	FUNGICIDE	MOA	AMOUNT PER ACRE	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
Seedling Disease	<i>azoxystrobin</i> Abound 2.08F	11	6 fl oz		Abound is active against Rhizoctonia damping off and Aspergillus crown rot.
Southern Stem Rot (White Mold) and Rhizoctonia Limb Rot	<i>azoxystrobin</i> Abound 2.08F	11	18.5-24.6 fl oz 12.3 fl oz (dryland only) These rates are for soilborne diseases to include suppression of CBR and for leaf spot diseases. Note: There is some evidence that the efficacy of <i>strobilurin</i> fungicides for management of leaf spot is not as strong as it once was. Therefore, growers should consider tank-mixing additional fungicides for management of leaf spot when using a <i>strobilurin</i> fungicide.	4 H/ 14 D	For increased disease control, growers should consider tank-mixing Alto (<i>cyproconazole</i> , 5.5 fl oz/A) with Abound (12-24.5 fl oz/A) 60-90 days after planting. Apply Abound, AzoxyStar, Artisan (26-32 fl oz/A), Evito, or Convoy (26 fl oz/A) as a foliar spray at 50-60 days after planting and reapply 28 days later. Note: Convoy and Artisan may also be applied on a 4-spray program. Consult label for more information. Begin treating immediately if active white mold is observed before the first scheduled application. Consult label for application strategies concerning the use of each product. Note: A fungicide to control leafspot must always be mixed with Convoy. Additional leaf spot control may not be needed if Artisan is used twice in a season at 26-32 fl oz/A (but tank-mixing and additional fungicide may be a good idea); however, if Artisan is applied in four applications, additional fungicides such as <i>chlorothalonil</i> or <i>thiophanate methyl</i> (e.g. Topsin M) must be added to supplement control of leaf spot. Maximum rate is 49 fl oz/A/season. Due to regulation from the European Union, current use of any product that contains <i>propiconazole</i> , such as Artisan, may be problematic. Growers should seek additional information before using such products.
	<i>azoxystrobin</i> AzoxyStar	11	12-24.5 fl oz see note	4H/ 14D	Do not apply more than 49 fl oz/A/season. Under heavy disease pressure and/or where there is high rainfall and/or irrigation, use 18.5-24.5 fl oz/A. For light disease pressure or dry environmental conditions, (non-irrigated, low rainfall) use 12-24.5 fl oz/A. Note: There is some evidence that the efficacy of <i>strobilurin</i> fungicides for management of leaf spot is not as strong as it once was. Therefore, growers should consider tank-mixing additional fungicides for management of leaf spot when using a <i>strobilurin</i> fungicide.
	<i>azoxystrobin</i> + <i>benzobendiflupyr</i> (solatenol) Elatus	11 + 7	0.5-0.65 fl oz/1000 linear feet 7.3-9.5 fl oz/A Elatus is effective against soilborne and leafspot diseases.	12 H/ 30 D	For suppression of stem rot/white mold, CBR and other diseases. Apply in a 7-10" band 14-21 days after planting. For foliar diseases begin 30-40 days after planting. Apply 7.3 fl oz/A on a 14-day schedule, or 9.5 fl oz/A on a 21-28 day interval. For soilborne diseases apply 7.3 fl oz/A 3 times on a 14-day interval, or 9.5 fl oz/A 2 times on a 21-28 day interval beginning 45-60 days after planting. Do not apply more than 21.9 fl oz/A per season
	<i>azoxystrobin</i> + <i>flutriafol</i> TopGuard EQ	11 + 3	8 fl oz	12 H/ 14 D	Note: TopGuard EQ, 5-7 fl oz/A, can be used for management of foliar diseases of peanut.
	<i>azoxystrobin</i> + <i>tebuconazole</i> HELMSTAR Plus SC	11 + 3	13 fl oz (approximately 60 and 90 days after planting)	12 H/ 14 D	Do not apply more than 51.6 fl oz/A in one season.
	<i>bixafen</i> + <i>flutriafol</i> LUCENTO	7 + 3	3-5 fl oz	12H/ 14D	For resistance management, do not make more than 2 sequential applications per year before alternating to another fungicide with a mode of action different than Group 7 or Group 3. Do not apply more than 11 fl oz of product per year.
	<i>flutolanil</i> Convoy	7	2 pt (1 application) 1-2 pt (2 applications) 0.5-1 pt (4 applications)	12 H/ 14 D	Convoy has no activity against leaf spot diseases. Therefore, a full-rate of an additional fungicide which is effective against leaf spot diseases should be tank-mixed with Convoy at each application. Maximum rate is 64 fl oz/A/season.

PEANUT DISEASE CONTROL

PEST	FUNGICIDE	MOA	AMOUNT PER ACRE	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS	
Southern Stem Rot (White Mold) and Rhizoctonia Limb Rot <i>(continued)</i>	<i>flutolanil + propiconazole</i> Artisan	7 + 3	26 or 32 fl oz (2 applications) 13 or 16 fl oz (4 applications)	12 H/ 40 D	The <i>propiconazole</i> in Artisan is an effective fungicide for the control of leaf spot diseases; however there are concerns for the potential of resistance to occur. When Artisan is applied at the 13-16 fl oz/A rates, the amount of <i>propiconazole</i> is not sufficient to control leaf spot and additional fungicide should be tank-mixed to improve efficacy. At the 26-32 fl oz/A rates of Artisan, the amount of <i>propiconazole</i> may be sufficient; however growers are advised to add additional fungicide to ensure adequate leaf spot protection. Maximum rate is 84 fl oz/A/season. NOTE: Because of issues with <i>propiconazole</i> and the European Union, growers are not encouraged to use Artisan at this time.	
	<i>fluxapyroxad + pyraclostrobin</i> Priaxor	7 + 11	8 fl oz	12 H/ 14 D	Priaxor has proven to be a very good replacement product for Headline in peanut disease management programs. Priaxor at a rate of 4 fl oz/A is appropriate for management of leaf spot; the 8 fl oz rate is appropriate for management of leaf spot and soilborne diseases. Maximum rate is 24 fl oz/A/season.	
	<i>fluoxastrobin</i> Evito 480 SC	11	5.7 fl oz	12 H/ 14 D	Evito 480 SC is in the same chemical class as Abound, Headline, and Stratego and should not be used in the same program with these products. Maximum rate is 22.8 fl oz/A/season.	
	<i>fluoxastrobin + tebuconazole</i> Evito T	3 + 11	9-11.2 fl oz	12 H/ 14 D	For control of white mold (stem rot), Rhizoctonia limb rot, and foliar diseases of peanut. Maximum rate is 44.8 fl oz/A/season.	
	<i>metconazole</i> Quash	3	2.5-4 fl oz	12 H/ 14 D		
	<i>penhopyrad</i> Fontelis	7	12-24 fl oz	12 H/ 14 D	Fontelis is a new fungicide in the SDHI class for disease management in peanut and will work well with <i>triazole</i> and <i>strobilurin</i> fungicides for resistance management. Make no more than 3 sequential applications before switching to a fungicide with a different mode of action. Maximum rate is 72 fl oz/A/season.	
	<i>pyraclostrobin</i> Headline	11	12-15 fl oz	12 H/ 14 D	Headline at this rate is an effective component of a soilborne program that also includes use of Provost, <i>tebuconazole</i> , Convoy, Quash, or Artisan. Note: There is recent concern that <i>strobilurin</i> fungicides are now less effective against leaf spot diseases and growers may consider tank-mixing additional fungicides with <i>strobilurin</i> fungicides for added protection from leaf spot. Maximum rate is 45 fl oz/A/season.	
	Labeled formulations of tebuconazole include, among others:					Apply <i>tebuconazole</i> products, or Quash (<i>metconazole</i>) 4 times per season starting at the second or third leaf spot spray. Growers are typically advised to tank-mix an additional leaf spot fungicide with <i>tebuconazole</i> products. Growers may also wish to tank-mix a fungicide with Quash for additional leaf spot control where leaf spot resistance to <i>triazole</i> fungicides has occurred. Maximum rate is 16.02 fl oz/A/season. The lower rate is generally considered effective for low-to-moderate pressure from white mold (stem rot) and Rhizoctonia limb rot. The higher rate is appropriate for more severe pressure from these diseases. Note: Research has shown that <i>tebuconazole</i> is often less effective in control of leaf spot than in the past. Where leaf spot is likely to be a problem, growers may consider tank-mixing <i>tebuconazole</i> with 0.75-1 pt/A <i>chorothalonil</i> or 5 fl oz Topsin (1st and 3rd applications only) to ensure adequate leaf spot control.
	<i>tebuconazole</i> Orius 3.6F, TriSum 3.6F, Integral 3.6F, T ebustar 3.6F, Muscle 3.6F, Tebuzol 3.6F, many others	3	7.2 fl oz Maximum rate is 28.8 fl oz/A/ season, except for Muscle 3.6 F (16 fl oz)	12 H/ 14 D		
	<i>tebuconazole + chlorothalonil</i> Muscle ADV	3 + M05	2 pt	12 H/ 14 D	Muscle ADV is a pre-mix of <i>tebuconazole</i> and <i>chlorothalonil</i> . Growers should note that the rate of Muscle ADV is 2 pt/A.	
<i>tebuconazole + prothioconazole</i> Provost Silver	3	11-13 fl oz	12 H/ 14 D	Provost is a new fungicide labeled for the control of leaf spot diseases, peanut rust, white mold, and Rhizoctonia limb rot. It is also labeled for suppression of CBR. Provost is applied in a 4-block program. Higher rates can be used in fields where disease is a particular problem. Maximum rate is 52 fl oz/A/season.		

PEANUT DISEASE CONTROL

PEST	FUNGICIDE	MOA	AMOUNT PER ACRE	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
Early emergence applications for management of White Mold and other Soilborne Diseases	<i>azoxystrobin</i> Abound 2.08F	11	0.4-0.8 fl oz/1000-row feet	4 H/ 14 D	Banded at 100% emergence for suppression of white mold. For greatest efficacy, concentrate the entire broadcast rate into a band wide enough to cover the peanuts. Check with your local Cooperative Extension office for recommendations on optimum timing of an early emergent application of Proline. Apply in a 4-6" band at 100% emergence. Abound: maximum rate is 49 fl oz/A/season. Proline: maximum rate is 22.8 fl oz/A/season.
	<i>prothioconazole</i> Proline 480SC	3	5.7 fl oz	24 H/ 14 D	
FOLIAR DISEASES					
Late leafspot (Cercosporidium personatum),	<i>cyproconazole</i> Alto 100SL	3	5.5 fl oz	12 H/ 30 D	See note for tetraconazole Domark 230ME.
Early leafspot (Cercospora arachidicola) and Rust (Puccinia arachidis)	<i>flutriafol</i> Rhyme	3	3.5-7 fl oz	12 H/ 7 D	
	TopGuard	3	7-14 fl oz		
	<i>fluoxastrobin + tebuconazole</i> Evito T	3	6-9 fl oz	24 H/ 14 D	Fungicide resistance management should be considered when this product is used with other <i>tebuconazole</i> products, Evito, Evito T, Stratego, Abound, Headline, Absolute, and Evito. Maximum rate is 44.8 fl z/A/season.
	<i>fluxapyroxad + pyraclostrobin</i> Priaxor	7 + 11	4-8 fl oz	12 H/ 14 D	Priaxor has proven to be a very good replacement product for Headline in peanut disease management programs. Priaxor at a rate of 4 fl oz/A is appropriate for management of leaf spot; the 6-8 fl oz rate is appropriate for management of leaf spot where the first fungicide application is made approximately 45 days after planting. Maximum rate is 24 fl oz/A/season.
	<i>picoxystrobin + cyproconazole</i> Approach Prima	11 + 3	5-6.8 fl oz	12 H/ 30 D	See note for pyraclostrobin Headline.
	<i>propiconazole + trifloxystrobin</i> Stratego	3	7 fl oz.	24 H/ 14 D	Fungicide resistance management should be considered when this product is used with <i>tebuconazole</i> products, Absolute, Evito, Evito T, Headline, or Abound. Maximum of 6 applications/season
	<i>pydiflumetofen</i> Miravis	7	3.4 fl oz	4 H/ 14 D	Apply on a 21-to-28 day interval. Note: Miravis is NOT effective against peanut rust disease.
	<i>pyraclostrobin</i> Headline	11	6-9 fl oz	24 H/ 12 D	NOTE: all soilborne fungicides described in the section "Control of Southern stem rot/white mold" area also active against leaf spot diseases EXCEPT for Convoy. Efficacy of these fungicides ranges from fair/good for <i>tebuconazole</i> to excellent for Provost and Headline. Most of the fungicides listed above fall into the "good" category. Headline (<i>pyraclostrobin</i>) is a <i>strobilurin</i> fungicide and is in the same chemical class as Abound, Evito, the fluoxastrobin component of Evito T, and the <i>trifloxystrobin</i> component of Stratego and Absolute. To best adhere to fungicide resistance management guidelines, Headline should not be used in the same program with these fungicides. Maximum rate is 45 fl oz/A/season.
	<i>tetraconazole</i> Domark 230ME	3	5.25-6.9 fl oz	12 H/ 14 D	NOTE: For foliar disease control, apply Alto 100SL up to two times as a part of a season-long disease management program; especially as a tank-mix partner for Abound. Alto maximum rate is 11 fl oz/A/season. Domark 230ME (2.5 fl oz/A) can be applied only if tank mixed with <i>chlorotalonil</i> (16.0 fl oz/A) or other leaf spot material. Do not apply more than 13.8 fl oz/A/ of Domark per season.
	<i>tetraconazole + chlorothalonil</i> Mazinga ADV	3 + M5	2 pt	12 H/ 14 D	Note: Rate of Mazinga ADV is 2 pt/A

PEANUT DISEASE CONTROL

PEST	FUNGICIDE	MOA	AMOUNT PER ACRE	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
FOLIAR DISEASES (continued)					
Late leafspot (Cercosporidium personatum), <i>(continued)</i>	<i>tetraconazole + thiophanate methyl</i> Acropolis	3 + 1	23 fl oz	24 H/ 14 D	Do not make more than four applications of this product per year.
	<i>tebuconazole + trifloxystrobin</i> Absolute	3 + 11	3.5 fl oz	12 H/ 14 D	Absolute is a combination of <i>tebuconazole</i> and <i>trifloxystrobin</i> . Fungicide resistance management should be considered when this product is used with <i>tebuconazole</i> products, Stratego, or Abound. Maximum of 4 applications/season.
Foliar Diseases	<i>chlorothalonil + sulfur</i> Bravo S is a pre-mix combination		68 fl oz		
	<i>chlorothalonil</i> Bravo Ultrex Bravo Weather Stik Chemnut 720 Chlorothalonil 720 Terranil 6L Echo 90DF Echo 720 Equus 720 Equus DF GK-Aragold 720	M5	1.36 lb 1.5 pt 1.5 pt 1.5 pt 1.5 pt 1.25 lb 1.5 pt 1.5 pt 1.36 lb 1.5 pt	12 H/ 14 D Except for Bravo S (24 H)	Apply <i>chlorothalonil</i> or <i>chlorothalonil + copper</i> on a 10-14 day interval. The exact interval between applications depends on rotation, weather, etc. Do not feed peanut hay treated with <i>chlorothalonil</i> to livestock. If rust is found in a field, and the peanuts are more than 3 weeks from expected harvest, apply <i>chlorothalonil</i> every 10 days until 2 weeks from harvest. If peanuts are 2 weeks or less from harvest, no control is necessary.
	Tank-mix combinations of <i>chlorothalonil</i> and <i>copper hydroxide</i> <i>chlorothalonil + Kocide 4.5LF</i>	M5	(0.75 lb ai/A) 1 pt		Do not mix any <i>copper</i> fungicide with Folicur.
	Tank-mix combinations of <i>chlorothalonil</i> Bravo + Tilt <i>propiconazole chlorothalonil + Tilt</i>	M5, 3	(0.75 lb ai/A) 2 oz	12 H/ 14 D	The rate of <i>chlorothalonil</i> when tank mixed with Tilt will vary depending on formulation (ie. 6 lb/gal would require 16 oz and 4.2 3 lb/gal would require 22.6 oz) If RUST is detected and peanuts are more than 3 weeks from expected harvest, use a full rate of <i>chlorothalonil</i> every 10 days until 2 weeks from harvest. Using DMIs (ie. Folicur and Tilt) full season increases the possibility for fungal resistance. If DMIs are used full season make the last leaf spot application with <i>chlorothalonil</i>. Tilt does not control rust. NOTE: Due to current considerations with the European Union, use of any product containing <i>propiconazole</i> is not recommended on peanuts. Maximum rate is 12 pints/A/season.
	<i>chlorothalonil + propiconazole</i> Echo-PropiMax	M5, 3	(0.75 lb ai/A) 2 oz	12 H/ 14 D	
	<i>chlorothalonil + propiconazole</i> Tilt/Bravo	M5, 3	1.5 pt	12 H/ 14 D	
	<i>thiophanate methyl</i> Topsin M 70WP T-Methyl 70W Topsin 4.5FL Topsin 4.5 FL + <i>tebuconazole</i>	1	1/2 lb 1/2 lb 10 fl oz 5 fl oz + 7.2 fl oz	24 H/ 14 D	Neither Topsin, <i>thiophanate methyl</i> 85WDG nor any other formulation of <i>thiophanate methyl</i> should be used alone for control of foliar diseases of peanut, but should be tank mixed and/ or rotated with other non-benzimidazole fungicides such as <i>chlorothalonil</i> . Use of <i>thiophanate methyl</i> should be restricted to a single solo application or 2 tank-mix applications per season. The maximum rate of <i>thiophanate methyl</i> 85WDG is 1.65 lb/A/season.
	Thiophanate Methyl 85WDG	1	0.4 lb	24 H/ 14 D	The rate for ELAST 400F is 12.8 fl oz if it is tank-mixed with another fungicide, e.g. <i>tebuconazole</i> for added control of peanut leaf spot. The rate is 15 fl oz if it is used alone or tank-mixed with a fungicide that is not effective against leaf spot.
	<i>dodine</i> ELAST 400F	U12	12.8-15 fl oz	48 H/ 14 D	
	<i>mancozeb</i> Koverall	M3	1-2 lbs	24 H/ 14D	

PEANUT DISEASE CONTROL

PEST	FUNGICIDE	MOA	AMOUNT PER ACRE	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
FOLIAR DISEASES (continued)					
Cylindrocladium Black Rot (CBR)	<i>metam sodium</i> 42%	M3	10 gal/A		To be effective, the fumigant <i>metam sodium</i> must be applied very carefully. To avoid injury to the seed and the seedlings, the fumigant must be applied at least 14 days before planting to a depth of 8-10". <i>Metam sodium</i> should be applied only when the soil temperature is greater than 60°F and when the soil moisture is like it would be for suitable seed germination. It is critical to get a good seal on the chisel trace left after fumigation so that the <i>metam sodium</i> does not escape directly into the atmosphere. The rows must be marked so that seed can be planted directly above where the fumigant was applied. Growers who are using this treatment for the first time may want to consult with their local county Cooperative Extension agent.
	<i>prothioconazole</i> Proline 480SC	3	0.4 fl oz/1000 row ft 5.7 fl oz/A	48 H/ 14 D	Proline 480SC is applied in-furrow for the management of CBR. See label for rate information and additional application strategies for management of CBR and perhaps white mold (stem rot). Maximum rate is 22.8 fl oz/A/season.
	<i>tebuconazole</i> Abound 2.08F tebuconazole 3.6F Priaxor Headline Provost Fontelis Elatus	11 3 11 + 7 11 3 7 11 + 3	18.5-24.6 fl oz 7.2 fl oz 12-15 fl oz 8-10.7 fl oz 8 fl oz/A 16-24 fl oz 7.3-9.5 fl oz	12 H/ 14 D Except Headline – 4 H	Provost <i>tebuconazole</i> , Abound, Elatus, Fontelis, Priaxor, and Headline are labeled for the “suppression” of CBR. This means that they may have some limited benefit to the grower in the management of this disease. However neither is likely to result in significant reduction in CBR when compared to the benefits of <i>metam sodium</i> .

RECOMMENDATIONS SET FORTH BY THE NORTH AMERICAN FUNGICIDE RESISTANCE ACTION COMMITTEE

1. Reduce initial inoculum (fungal populations) through good cultural practices.
2. Do not use less than the minimum label rate alone or in tank mixtures.
3. If more than four sprays of DMI (for example *tebuconazole* and *propiconazole*) fungicides will be made in a season, it is strongly recommended that all DMI sprays be mixed with an effective non-DMI fungicide.
4. Use in a preventative application schedule.
5. Calibrate sprayer and configure spray tips to ensure thorough coverage of peanut foliage.
6. DMI fungicides are not recommended for season-long use alone. Use alternating blocks of sprays of DMI fungicides with non-DMI fungicides, OR use tank mixes of DMI and non-DMI fungicides. See label directions!
7. Alternating sprays or tank mixtures with other DMI fungicides will not help prevent resistance development.
8. When using a *strobilurin* fungicide as a solo product, for example Headline, Abound, or Evito, the number of applications should be no more than 1/3 (33%) of the total number of fungicide applications per season.
9. For *strobilurin* mixes (e.g. Absolute, Evito T and Stratego) in programs which tank mixes or pre-mixes a *strobilurin* with mixing partners of a different mode of action are used, the number of *strobilurin* containing applications should be no more than 1/2 (50%) of the total number of fungicide applications per season.
10. In programs in which applications of *strobilurin* fungicides are made with both solo products and mixtures, the number of *strobilurin* containing applications should be no more than 1/2 (50%) of the total number of fungicide applications per season.

PEANUT SEED TREATMENT

Bob Kemerait, Extension Plant Pathologist, Tim Brenneman and Albert Culbreath, Plant Pathologists

Dynasty PD (<i>azoxystrobin + fludioxonil + mefenoxam</i>)	3-4 fl oz/100 lb
Vitavax PC	4-5 oz/100 lb
Rancona VPD (<i>ipconazole + carboxin + metalaxyl</i>)	4 oz/100 lb seed

PEANUT NEMATODE CONTROL

PEST	NEMATICIDE	AMOUNT PER ACRE	REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
Nematodes	<i>fluopyram + imidacloprid</i> Velum Total	18 fl oz	12 H/ 14 D	Apply specified dosage in the following methods: 1) In-furrow spray during planting directed on or below seed; 2) Chemigation into the root-zone through low pressure drip or trickle irrigation. Do not apply more than 19 fl oz/A of Velum Total per year. Do not apply Velum Total within 30 days of harvest. Regardless of formulation or method of application, apply no more than 0.5 lb <i>imidicloprid</i> or 0.45 lb <i>fluopyram ai/A/year</i> , including seed treatment, soil and foliar uses. Note: PROPULSE (<i>fluopyram + prothioconazole</i>) can be applied at 13.6 fl oz/A at pegging time for additional control of nematodes following an in-furrow application of Velum Total. Propulse offers additional nematode control through chemigation that uses overhead irrigation equipment. Applications should be made using 0.1 to 0.25 acre-inch of water to move PROPULSE into the soil.
	Aldicarb AgLogic 15G	7 lb/A at-plant followed by 10 lb/A post emergence.		At-Planting: Apply granules in a 4-6" band over open seed furrow (T-band) and immediately cover with soil by mechanical means. OR Apply granules in a 6-12" band and immediately cover with soil by mechanical means to a depth of 2-4". Plant seed into treated zone. Post-Emergence: Apply granules in a band 12-18" wide on the row and into the plant canopy. Ensure that plant foliage is dry prior to application. Dislodge granules from foliage by suitable means that will not damage the plant. Apply 6-9 gal/A rates broadcast with a moldboard plow. Apply 4.5-6 gal/A rates with a single chisel in-row application. All applications should be made at least 7-14 days before planting.
	Telone II	4.5-9 gal	5 D Post application/ -	Apply 6-9 gal/A rates broadcast with a moldboard plow. Apply 4.5-6 gal/A rates with a single chisel in-row application. All applications should be made at least 7-14 days before planting.
	Vydate C-LV	17-34 fl oz	48 H/ -	Vydate C-LV is used to supplement use of a pre-plant or at-plant application of a nematicide. (If Vydate is used at-plant, it should be applied at 34.0 fl oz/A.) Recommended foliar application is a split application of 17 fl oz/A at 14-28 days after planting followed by a second application of 17 fl oz/A 14 days later. Best used in low nematode fields. Maximum rate is 136 fl oz/A/season.
A nematicide application at pegging is recommended regardless of which nematicide is used at planting. DO NOT FEED PEANUT HAY TREATED WITH THESE NEMATICIDES TO LIVESTOCK.				

NEMATICIDE CONTROL RATING

NEMATICIDE	RATING
Telone II	excellent
Vydate	poor to fair

Nematicides listed below are rated on a scale of excellent, good, fair and poor. Those listed as poor to fair would be acceptable in low nematode population fields, but should not be used in fields where root-knot nematodes cause severe problems.

PEANUT WEED CONTROL

Eric P. Prostko, Extension Agronomist – Weed Science

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION/A	LBS ACTIVE INGREDIENT/A		
EARLY PRE-PLANT FOLIAR BURNDOWN OF EMERGED ANNUAL WEEDS AND/OR COVER CROPS IN REDUCED TILLAGE SYSTEMS					
<i>glyphosate</i> various trade names 3 lb ae/gal 3.73 lb ae/gal 4 lb ae/gal 4.17 lb ae/gal 4.50 lb ae/gal 5 lb ae/gal	9	16-48 oz 13-39 oz 12-36 oz 11.7-35 oz 11-32 oz 10-29 oz	0.38-1.13 ae	4 H/ –	Apply any time prior to planting to control emerged weeds. Refer to specific label for weeds controlled, application rates, adjuvants, and precautions. <i>Glyphosate</i> does not adequately control cutleaf eveningprimrose or Carolina geranium, and may not provide acceptable control of wild radish. For cover crop control only, use the following rates: wheat < 12”, 0.56 lb ae/A; wheat > 12”, 0.75 lb ae/A; rye < 18”, 0.56 lb ae/A; rye > 18”, 0.75 lb ae/A. <i>Glyphosate</i> can also be tank-mixed with Valor (1-3 oz/A), Aim (1-2 oz/A), or ET (0.5-2 oz/A) to improve the spectrum of control, particularly for annual morningglories. Refer to specific comments for Valor. Sequence 5.25EC (<i>glyphosate</i> + <i>S-metolachlor</i>) is also labeled for pre-plant use in peanut at 2.5-3.4 pt/A. Sequence at 3.4 pt/A is equivalent to 1.28 lb ai/A of <i>S-metolachlor</i> + 0.96 lb ai/A <i>glyphosate</i> . Applications of <i>glyphosate</i> to wheat and rye should be made before the boot stage or after the small grain is fully headed.
<i>paraquat</i> various trade names 2SL 3SL	22	40-60 oz 27-40 oz	0.63-0.94	12 H/ –	Apply prior to planting to control emerged weeds. Add non-ionic surfactant at 1qt/100 gals or crop oil at 1 gal/100 gals. Paraquat will not adequately control horseweed, swinecress, purslane speedwell, curly dock, cutleaf eveningprimrose, and larger wild radish. For cover crop control only, use the following rates: wheat, 0.63 lb ai/A (2.5 pt/A of 2 lb/gal or 1.7 pt/A of 3 lb/gal); rye, 0.50 lb ai/A (2 pt/A of 2 lb/gal or 1.3 pt/A of 3 lb/gal). Cover crops must be mature (seedheads) for adequate control. Can also be tank-mixed with Valor (1-3 oz/A) to improve the spectrum of control and provide residual weed control. Refer to specific comments for Valor. **All applicators of paraquat must complete the EPA’s training program before mixing, loading and applying Refer to the following web-site for more information: https://www.epa.gov/pesticide-worker-safety/paraquat-dichloride-training-certified-applicators
<i>2,4-D amine</i> various trade names 3.8 lb/gal	4	8-16 oz	0.24-0.48	48 H/ –	Tank-mix with <i>glyphosate</i> or <i>paraquat</i> . <i>2,4-D</i> is the most cost-effective option available for burndown of cutleaf eveningprimrose. <i>2,4-D</i> does not control Carolina geranium. Some <i>2,4-D</i> products are labeled for application to previous crop stubble or fallow land. In this case, the label directs the user to not plant a crop “until 3 months after application or until the product disappears from the soil.” UGA research suggests that peanut can be planted 7 days after a pre-plant burndown application of <i>2,4-D</i> .
<i>thifensulfuron + tribenuron</i> FirstShot 50SG	2	0.5-0.8 oz	0.008-0.013 + 0.008-0.013	12 H/ –	Tank-mixed with <i>glyphosate</i> or <i>paraquat</i> . FirstShot will help improve the control of many broadleaf weeds such as henbit, wild radish, Carolina geranium, and chickweed. Peanut can be planted 30 days after application. Add a NIS at 0.25% v/v or COC at 1% v/v unless tank-mixed with a loaded <i>glyphosate</i> formulation.
<i>carfentrazone</i> Aim 2EC	14	1-2 oz	0.016-0.031	12 H/ 7 D	Tank-mix with <i>glyphosate</i> to improve the burndown control of annual morningglory, tropical spiderwort, and small pigweed (< 1” tall). Apply prior to planting or up until 24 hours after planting. Add a NIS at 0.25% v/v or COC at 1% v/v unless tank-mixed with a “loaded” <i>glyphosate</i> formulation.
<i>pyraflufen</i> ET 0.208EC	14	0.5-2 oz	0.001-0.003	12 H/ 7 D	Tank-mix with <i>glyphosate</i> to improve the burndown control of annual morningglory and small pigweed (< 1” tall). Apply as pre-plant burndown or after planting but before crop emergence. Add a COC at 1% v/v unless tank-mixing with a “loaded” <i>glyphosate</i> formulation.

PEANUT WEED CONTROL

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION/A	LBS ACTIVE INGREDIENT/A		
PRE-PLANT SOIL INCORPORATED					
<i>ethalfuralin</i> Sonalan HFP 3EC	3	32 oz	0.75	24 H/ –	Controls annual grasses and small-seeded broadleaf weeds. Soil incorporate 2-3 inches deep within 2 days of application. Incorporation with implements other than power tiller requires two passes, preferably at cross angles. May be tank-mixed with Frontier/Outlook or Dual for control of mixed infestations of annual grasses and nutsedge. Sonalan may also be applied as a surface application to freshly prepared seedbeds but must be incorporated by 0.5-1” of rainfall or irrigation within 2 days after application.
<i>pendimethalin</i> Prowl/Pendimax 3.3EC Prowl H ₂ O 3.8 ACS	3	29-38 oz 32 oz	0.75-1 0.95	24 H/ –	Controls annual grasses and small-seeded broadleaf weeds. Soil incorporate 1-2 inches deep within 7 days of application. Incorporation with implements other than power tiller requires two passes, preferably at cross angles. Use high rate for Texas panicum or where heavy weed populations are anticipated. May be tank-mixed with Frontier/Outlook, Dual, or Pursuit for control of mixed infestations of annual grasses and nutsedge. Prowl can be applied immediately after planting to a freshly prepared seedbed up to 2 days after planting but before crop emergence. However, adequate incorporation in the form of 0.75” of irrigation or rainfall is needed within 48 hours for optimum activation when applied by this method. In strip-tillage production systems with heavy cover, the rate of <i>pendimethalin</i> should be increased to 42 oz/A (Prowl 3.3EC) or 35 oz/A (Prowl H ₂ O).
<i>metolachlor</i> Stalwart, Parallel PCS, Me-Too Lachlor 8EC <i>S-metolachlor</i> Dual Magnum 7.62EC Dual II Magnum 7.64EC Cinch 7.62EC EverpreX 7.62EC	15	16 oz 16 oz	1 0.95	24 H/ 90 D	Controls some annual grasses (not Texas panicum) and small-seeded broadleaf weeds. May provide limited Florida beggarweed suppression. Controls or suppresses yellow nutsedge but not purple nutsedge. Incorporation with implements other than power tiller requires two passes, preferably at cross angles. Deep incorporation may reduce effectiveness. May be tank-mixed with Prowl/Pendimax or Sonalan to control mixed infestations of annual grasses and yellow nutsedge. PPI treatments generally provide better control of nutsedge. Heavy rainfall after planting and/or non-uniform incorporation may result in crop injury expressed as delayed emergence and stunted growth of emerging plants. The generic formulations of <i>metolachlor</i> (Parallel PCS, Stalwart, Me-Too-Lachlor) have not provided the same length of residual control of certain weeds as similar rates of Dual Magnum formulations in some UGA field trials.
<i>diclosulam</i> Strongarm 84WG	2	0.45 oz	0.024	12 H/ –	Provides general broadleaf weed control. Incorporate into top 1-3” of final seedbed. Good to excellent control of many species including bristly starbur, wild poinsettia, eclipta, and copperleaf. Should be tank-mixed with a grass herbicide. Poor control of sicklepod. Control of nutsedge has been variable and inconsistent. Can also be applied pre-emergence. Crop rotation restrictions: soybeans – 0 months; wheat, barley – 4 months; oats, rye – 6 months; cotton – 10 months; corn, sorghum, tobacco – 18 months; other crops – 30 months. Pre-slurry in water before adding to larger spray/mix tank.
<i>imazethapyr</i> Pursuit 2AS 70DG	2	4 oz 1.44 oz	0.063	4 H/ 85 D	Controls purple and yellow nutsedge, wild poinsettia, wild radish, pigweed, burgherkin, and several other annual species. Does not control Florida beggarweed or sicklepod. Shallow incorporation is preferred. May be tank-mixed with Dual, Prowl/Pendimax, or Sonalan. Incorporated treatments are more persistent than pre-emergence or post-emergence applications and are more likely to result in carryover. Rotation intervals for various crops include the following: peanut, peas, southern pea, soybean – anytime; snap bean – 2 months; rye and wheat – 4 months; field corn – 8.5 months; barley, tobacco – 9.5 months; cotton, oat, sorghum, sunflower, sweet corn – 18 months.

PEANUT WEED CONTROL

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION/A	LBS ACTIVE INGREDIENT/A		
CHEMIGATION					
<i>metolachlor</i> Stalwart, Parallel PCS Me-Too-Lachlor	15	Refer to PPI section for rates			May be applied by injection through center pivot irrigation systems. Use at normal recommended rates. Apply after planting but before crop emergence. Requires proper system calibration and safety devices (check valves, cutoff switches, etc.) to provide effective weed control and prevent environmental contamination. Accurate herbicide application through chemigation may provide superior weed control compared to conventional ground applications. The generic formulations of <i>metolachlor</i> (Parallel PCS, Stalwart, Me-Too-Lachlor) have not provided the same length of residual control of certain weeds as similar rates of Dual Magnum formulations in some UGA field trials.
<i>S-metolachlor</i> Dual Magnum 7.62E Cinch 7.64EC	15	Refer to PPI section for rates			
<i>pendimethalin</i> Prowl/Pendimax 3.3EC Prowl H ₂ O 3.8ACS	3	Refer to PPI section for rates			
PRE-EMERGENCE					
<i>imazethapyr</i> Pursuit 2AS Pursuit 70DG	2	4 oz 1.44 oz	0.063	4 H/ 85 D	See comments for Pursuit PPI. Controls the same weeds as listed for Pursuit PPI but with greater dependency on rainfall or irrigation for activation.
<i>metolachlor</i> Stalwart Parallel PCS Me-Too-Lachlor	15	16 oz	1-1.33	24 H/ 90 D	Controls some annual grasses (not Texas panicum) and small-seeded broadleaf weeds. Provides some suppression of sicklepod and Florida beggarweed. Apply after planting and before crop and weeds emerge. If Dual is used as a PPI treatment, any additional application of Dual should be delayed until peanuts begin emerging (AC). Multiple applications – pre-plant incorporated followed by at-cracking treatments – improve control of sicklepod, Florida beggarweed, and yellow nutsedge. Pre-emergence treatments generally provide better broadleaf weed control/suppression Do not apply more than 2.66 pt/A/year of Stalwart/Parallel/Me-Too-Lachlor or 2.8 pt/A/ year of Dual Magnum/Dual II Magnum/Cinch formulation. The generic formulations of <i>metolachlor</i> (Parallel, Stalwart, Me-Too-Lachlor) have not provided the same length of residual control of certain weeds as similar rates of Dual Magnum formulations in some UGA field trials. PPI or PRE applications of Dual should be avoided if excessive rainfall is anticipated.
<i>S-metolachlor</i> Dual Magnum 7.62EC Dual II Magnum 7.64EC Cinch 7.64EC		16 oz	0.96-1.27		
<i>diclosulam</i> Strongarm 84WG	2	0.45 oz	0.024	12 H/ –	Refer to PPI section.
<i>flumioxazin</i> Valor SX 51WG Valor EZ 4SC Panther 51WG Panther 4SC Rowel 51WG Red Eagle Flumioxazin 51WG	14	3 oz	0.096	12 H/ –	Apply immediately after planting but no later than 2 days after planting. Plant peanuts at least 1.5” deep. DO NOT irrigate when peanuts are cracking. Rainfall or irrigation at cracking will cause temporary crop injury that should not result in reduced yields if applied according to the label. Valor will provide good to excellent control of many broadleaf weeds including Florida beggarweed, Palmer amaranth, and tropic croton. Valor will not control annual/perennial grasses, sicklepod, nutsedge, and cocklebur. Valor can be tank-mixed with Prowl, Sonalan, Dual Magnum, or Warrant. Dual or Warrant tank-mixes with Valor would only be suggested when planting in fields with a history of tropical spiderwort and/or yellow nutsedge. Can also be used in strip-tillage peanut production systems in combination with glyphosate or paraquat to improve burndown control. Rotation restrictions include the following: soybeans – 0 months; field corn – 1 month; cotton, tobacco, wheat – 2 months. Refer to current product label for additional rotational restrictions. Completely clean spray equipment THE SAME DAY OF USE as directed on the herbicide label! Pre-slurry all <i>flumioxazin</i> formulations in water before adding to larger spray/mix tank.

PEANUT WEED CONTROL

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION/A	LBS ACTIVE INGREDIENT/A		
PRE-EMERGENCE					
<i>sulfentrazone + carfentrazone</i> Spartan Charge 3.5L	14 + 4	3-3.75 oz	0.074-0.092 + 0.008-0.01	12 H/ –	Will provide good to excellent residual control of pigweed. Can be tank-mixed with Prowl or Dual. Can be tank-mixed with <i>glyphosate</i> or <i>paraquat</i> pre-plant burndown in strip-tillage systems. Can be applied up to 3 days after planting. However, do not apply after peanut emergence, at cracking, or if seedling is close to the soil surface. Do not use on soils classified as sand, which have less than 1% OM. Do not irrigate when peanuts are cracking. Rotation Restrictions: soybeans, sunflowers, tobacco – anytime; field corn, small grains – 4 months; sorghum – 10 months; cotton – 18 months; canola – 24 months.
<i>acetochlor</i> Warrant 3ME	15	48 oz	1.125	12 H/ Forage 90 D	Can be applied PRE and/or early POST (after crop emergence up through the R1 stage of growth. R1 ends when 50% of plants have a visible peg (R2)). Will provide good to excellent control of annual grasses (except Texas panicum), pigweed, and tropical spiderwort. Crop injury potential will increase if used in cold/wet soils. In wet years or environments, the length of residual control with Warrant may be reduced.
AT CRACKING OR EARLY POST-EMERGENCE					
<i>imazethapyr</i> Pursuit 2.0 AS 70 DG	2	4 oz 1.44 oz	0.063	4 H/ 95 D	See comments for Pursuit PPI and PRE. Provides effective control of nutsedge, wild poinsettia, wild radish, bristly starbur, prickly sida, and several other annual species. Weed size is especially critical for effective control of nutsedge, bristly starbur, and prickly sida. If weeds are emerged, surfactant or crop oil concentrate should be included. May be tank-mixed with <i>paraquat</i> or 2,4-DB for broader spectrum control of emerged weeds.
<i>metolachlor</i> Stalwart Parallel PCS Me-Too-Lachlor 8EC	15	16 oz	1-1.33	24 H/ 90 D	See comments for Dual PPI and PRE. Compared to PPI and PRE treatments, AC applications provide better control of non-emerged broadleaf weeds such as Florida beggarweed and sicklepod. May be tank-mixed with <i>paraquat</i> + Basagran, <i>paraquat</i> + Storm or Cadre or Cobra treatments for improved contact activity and for suppression/control of problem broadleaf weeds and yellow nutsedge. Do not use Dual II Magnum/Cinch formulations after peanut emergence. Do not apply more than 2.66 pt/A/year of Stalwart/Parallel/Me-Too-Lachlor or 2.8 pt/A/year of Dual Magnum. Research has shown that Dual will provide good to excellent residual control of tropical spiderwort if applied before weed emergence. The generic formulations of <i>metolachlor</i> (Parallel PCS, Stalwart, Me-Too-Lachlor) have not provided the same length of residual control of certain weeds as similar rates of Dual Magnum formulations in some UGA field trials. Adjuvants are not needed when Dual is tank-mixed with POST treatments.
<i>S-metolachlor</i> Dual Magnum 7.62EC (various others)		16 oz	0.95		
<i>acetochlor</i> Warrant 3ME	15	48 oz/A	1.125	12 H/ Forage 90 D	Can be applied after emergence through the R1 (flowering) stage of growth. R1 ends when 50% of plants have a visible peg (R2). Can be tank-mixed with <i>paraquat</i> + Basagran, or <i>paraquat</i> + Storm treatments or Cadre or Cobra. Total amount of Warrant that can be applied PRE + early POST = 6 pt/A/year. Use a NIS at 0.25% v/v when tank-mixed with POST herbicides.
<i>pyroxasulfone</i> Zidua 85WG Zidua 4.17SC	15	1.5-2.0 oz 2.45-3.25 oz	0.08-0.11	12 H/ 0 D	Apply post-emergence for the residual control of certain annual weed species, including Palmer amaranth, Florida beggarweed, and annual grasses. Tank-mix with <i>paraquat</i> , Cadre, Cobra, or Ultra Blazer. Can be applied from V2 stage thru beginning of pod development stage (R3). Do not apply pre-emergence to peanuts. A total of 2 applications (4 oz/A of 85WG or 6.5 oz/A of 4.17SC/year total) can be applied. When applied POST with other herbicides, use a NIS at 0.25% v/v. UGA weed scientists have collected more data with the 85WG formulation than the 4.17SC formulation.

PEANUT WEED CONTROL

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION/A	LBS ACTIVE INGREDIENT/A		
AT CRACKING OR EARLY POST-EMERGENCE (continued)					
<i>paraquat</i> various trade names 2SL 3SL	22	5.4 oz 8 oz	0.125	12 H/ –	Provides effective, broad-spectrum weed control. Provides some suppression of yellow nutsedge. Apply anytime up to 28 days after ground crack. Include NIS at 0.25% (1 qt/100 gal) with all <i>paraquat</i> treatments. The success of “at-crack” sprays can be greatly improved by: 1) applying herbicides in a minimum of 15 GPA; 2) using flat fan nozzles; 3) decreasing ground speed; and 4) using lower spray pressures (30 PSI). Research indicates no adverse effects of adding <i>chlorothalonil</i> products with <i>paraquat</i> tank-mixtures where fungicide treatments are needed.
<i>paraquat</i> 3 SL or 2SL + <i>bentazon</i> + <i>acifluorfen</i> Storm 4EC	22 + 6 + 14	8 oz 12 oz + 16 oz	0.188 + 0.33 + 0.17	48 H/ 75 D	*Warrant 3ME (48 oz/A) or Dual Magnum/generic <i>metolachlor</i> (16 oz/A) or Zidua (1.5 oz/A of 85WG or 2.45 oz/A of 4.17SC) can be used in combination with this treatment to provide residual control of pigweed and tropical spiderwort. NIS is not recommended if Dual Magnum or generic <i>metolachlor</i> is used with <i>paraquat</i> + Storm. **All applicators of paraquat must complete the EPA’s training program before mixing, loading and applying Refer to the following website for more information: https://www.epa.gov/pesticide-worker-safety/paraquat-dichloride-training-certified-applicators
<i>paraquat</i> 3 SL or 2SL + <i>bentazon</i> Basagran, Broadloom 4EC 5EC	22 + 6 + 14	8 oz 12 oz + 8-16 oz 6.4-12.8 oz	0.189 + 0.25 + 0.5	48 H/ Forage 50 D	Provides effective, broad-spectrum weed control. Provides some suppression of yellow nutsedge. Generally reduces peanut injury compared to other <i>paraquat</i> treatments. The lower rate of Basagran (0.5 pt) is usually sufficient to reduce peanut foliar burn and provide control of smallflower morningglory. The higher rate (1 pt) is necessary for control of weeds such as bristly starbur and prickly sida. Apply anytime up to 28 days after ground crack. Include NIS at 1 qt/100 gal spray solution with all <i>paraquat</i> treatments. The success of “at-crack” sprays can be greatly improved by 1) applying herbicides in a minimum of 15 GPA; 2) using flat fan nozzles; 3) decreasing ground speed; and 4) using lower spray pressures (30 PSI). Research indicates no adverse effects of adding <i>chlorothalonil</i> products with <i>paraquat</i> tank-mixtures where fungicide treatments are needed. *Warrant (48 oz/A) or Dual Magnum or generic <i>metolachlor</i> (16 oz/A) or Zidua (1.5 oz/A of 85WG or 2.5 oz/A of 4.17SC) can be used in combination with this treatment to provide residual control of pigweed and tropical spiderwort. NIS is not recommended if Dual Magnum or generic <i>metolachlor</i> is used with <i>paraquat</i> + Basagran.
<i>diclosulam</i> Strongarm 84WG	2	0.45 oz	0.024	12 H/ –	24(c) label for use in Georgia. The only weed on current 24(c) label is tropical spiderwort but Strongarm also has POST activity on annual morningglory, bristly starbur, common cocklebur, common ragweed, eclipta, and horseweed. Can be applied up until 30 days after planting. Use in combination with a NIS at 0.25% v/v (1 qt/100 gals). When applied post-emergence in peanut, cotton rotation restriction is 18 months. Follow other rotation restrictions listed in PPI section. Label must be in the possession of user at the time of application.

PEANUT WEED CONTROL

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION/A	LBS ACTIVE INGREDIENT/A		
POST-EMERGENCE					
<i>acifluorfen</i> Ultra Blazer 2L	14	16-24 oz	0.25-0.38	- / 75 D	Especially useful for control of morningglories, tropic croton, wild radish, wild poinsettia, hophornbeam copperleaf, and spider flower. Adjust rate according to weed size and species as noted on the label. Use 16 oz/A or less for control of highly sensitive species such as hemp sesbania and showy crotalaria. Slight to moderate peanut foliage burn may result. Do not apply more than 2 pt/A per season as a post-emergence treatment. Apply with nonionic surfactant at 1 qt/100 gal spray solution (0.25% v/v). May be tank-mixed with 2,4-DB (1 pt/A). The Blazer + 2,4-DB tank mixture is generally more injurious to peanuts than either product alone. May be tank-mixed with Basagran for improved control of broadleaf weeds such as cocklebur, and prickly sida. A pre-packaged mix of <i>acifluorfen</i> + <i>bentazon</i> is marketed as Storm. Ultra Blazer can be tank-mixed with Dual Magnum or Warrant or Zidua + 2,4-DB. Generic formulation of <i>acifluorfen</i> is sold as Avalanche Ultra (Winfield) but has not been tested by UGA. Rain-free period for Ultra Blazer is 4 hrs.
<i>bentazon</i> Basagran, Broadloom 4EC 5EC	6	24-32 oz 19-26 oz	0.75-1	48 H/ Forage-50 D	Apply for post-emergence control of yellow nutsedge, cocklebur, bristly starbur, smallflower morningglory, prickly sida, and certain other weeds. Treat when broadleaf weeds are small and actively growing. Adjust rate according to weed size as noted on label. Two applications will be required for control of yellow nutsedge. For yellow nutsedge, include crop oil concentrate at 1 qt/A. Do not foliarly apply <i>sulfur</i> 14 days before or after use of crop oil concentrate to minimize risk of peanut foliage burn. May be tank-mixed with 2,4-DB amine 2L (0.5 pt/A) for improved control of morningglories. Early-season applications of <i>bentazon</i> at high rates following in-furrow applications of Di-Syston may infrequently result in SEVERE peanut injury. Rain-free period for Basagran is 4 hours.
<i>bentazon</i> + <i>acifluorfen</i> Storm 4EC	6 + 14	1.5 pt	0.5 + 0.25	48 H/ 75 D	Controls morningglories, cocklebur, prickly sida, ragweed, eclipta, tropic croton, and several other broadleaf weeds with less injury than Blazer alone. Application timing is critical—weeds must be small. Include surfactant or crop oil concentrate. Can be mixed with 2,4-DB for control of larger weeds and for control of sicklepod. May be tank-mixed with <i>paraquat</i> . Rain-free period for Storm is 4 hours.
2,4-DB 1.75SL 2SL	4	14.4-18.3 oz 13-16 oz Rate depends on formulation. Check product labels.	0.20-0.25 0.20-0.25 0.22-0.38 0.22-0.40	48 H/ 60 D	Apply up to 2 applications per season as an over-the-top treatment for broadleaf weed control. Use rates and application timing varies by specific product label. Apply in the seedling stage for control of morningglory and citronmelon. Cocklebur that is 1 foot or more in height can be controlled; however, earlier treatment is preferred. Also effective for control of escaped sicklepod. Do not apply if peanuts are under drought stress. Research indicates no adverse effects of adding <i>chlorothalonil</i> products with 2,4-DB where fungicide treatments are needed. Rain-free period for 2,4-DB is 1 hour. Do not tank-mix with post-emergence grass herbicides. If 2,4-DB is tank-mixed with grass herbicides, such as Poast or Select, reduced control of grasses can occur (approximately 8-15% reduction in control about 50% of the time).
<i>imazethapyr</i> Pursuit 2 AS 70 DG	2	4 oz 1.44 oz	0.25-0.38	4 H/ 85 D	See comments for Pursuit PPI, PRE, and AC/EP. Generally should be used early post-emergence when weeds are extremely small. Controls wild radish, pigweeds, morningglories, cocklebur, and several other annual species. Compared to PPI, PRE, and AC/EP treatments, POST applications are less effective on nutsedge, wild poinsettia, and some other species. Applications should be made before nutsedge exceeds 3-4 inches and bristly starbur exceeds 2". May be tank-mixed with <i>paraquat</i> or 2,4-DB. Post control of escaped wild poinsettia is greatly enhanced in combination with <i>paraquat</i> . Rain-free period for Pursuit is 1 hour. If 2,4-DB is tank mixed with grass herbicides, such as Poast or Select, reduced control of grasses can occur (approximately 8-15% reduction in control about 50% of the time).

PEANUT WEED CONTROL

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION/A	LBS ACTIVE INGREDIENT/A		
POST-EMERGENCE (continued)					
<i>imazapic</i> Cadre/Impose 2AS	2	4 oz	0.063	12 H/ 90 D	Provides excellent control of many broadleaf and grass weeds and both purple and yellow nutsedge. Apply as an early post-emergence treatment when weeds are less than 2-3 inches in height. Under conditions of heavy weed pressure, applications of Cadre 10-14 days following an at-cracking treatment (<i>paraquat</i> combination) has resulted in superior weed control. Use with NIS (0.25% v/v) or COC (1 qt/A). <u>Tank-mixing Select or Poast with Cadre can result in a 10-25% reduction in grass control.</u> A better alternative to this tank-mix would be to apply the grass herbicide first, then apply Cadre 3 days later. Rotation restrictions include: wheat, rye – 4 months; corn, snapbean, southern peas, soybeans, tobacco – 9 months; cotton, oats, sweet corn, grain sorghum – 18 months; canola – 40 months. See label for additional restrictions. Rain-free period for Cadre is 3 hours. Cadre can be tank-mixed with Dual Magnum, Warrant, or Zidua + 2,4-DB.
<i>lactofen</i> Cobra 2EC	14	12.5 oz	0.195	12 H/ 45 D	Apply after peanuts reach 6 true leaf stage of growth. Use a crop oil concentrate at 1% v/v (1 gal/100 gals). Provides good control of pigweeds, morningglories, ragweed, copperleaf, wild poinsettia, and eclipta. Cobra can be tank-mixed with Basagran, Dual Magnum, Cadre, Pursuit, Select, Warrant, Zidua, and 2,4-DB. Rain-free period is 30 minutes.
<i>sethoxydim</i> Poast 1.5EC Poast Plus 1EC	1	16-24 oz 24-36 oz	0.19-0.28	12 H/ 40 D	For control of annual and perennial grasses. Apply when annual grasses are small (1-6 inches) and actively growing. Under favorable conditions, large Texas panicum can be controlled. For perennial grass control, two applications are usually required for satisfactory control. Apply with a COC at 1% v/v (1 gal/100 gal). Tank-mixtures with other herbicides, such as 2,4-DB, may reduce grass control. Do not apply <i>sulfur</i> 14 days before or after application to minimize risk of peanut foliage burn. Reduced spray volumes (10 GPA) may improve grass control. Refer to Table 1 at the end of this chapter for more specific information about grass control. Rain-free period for Poast is 1 hour.
<i>clethodim</i> Select, Arrow, others 2EC Select Max/TapOut 0.97EC Section Three, Shadow 3EC	1	6-8 oz 12-16 oz 3.84-5.33 oz	0.09-0.125	24 H/ 40 D	For control of annual and perennial grasses. Apply when grasses are small (< 6 inches) and actively growing. Under favorable conditions, large Texas panicum and bermudagrass can be effectively controlled. Heavy bermudagrass pressure or larger Texas panicum requires a follow-up treatment. When tank-mixing with a broadleaf herbicide or controlling perennial grasses, increase rates (8-16 ozs/A-Select; 16-32 oz/A-Select Max/TapOut). Do not apply more than 21.33 oz/A/year (Section Three, Shadow), 32 oz/A/year (Select) or 64 oz/A/year (Select Max/TapOut). Always apply with a crop oil concentrate at 1% v/v (Select/Arrow). A NIS (0.25% v/v) can be used with Select Max/TapOut to reduce crop injury potential. May be tank-mixed with Basagran, Blazer, Storm, or Orthene. Do not tank-mix with chlorothalonil products or reduced grass control can occur. Refer to Table 1 at the end of this chapter for more specific information about grass control. Rain-free period is 1 hour.
<i>fluzifop-P</i> Fusilade DX 2EC	1	8-24 oz	0.125-0.375	12 H/ 40 D	For the control of annual and perennial grass weeds. Use rate depends on weed and weed size. Refer to table at the end of this section for specific information about rates and timings. Do not apply more than 48 oz/A/season. Do not apply more than 24 oz/A/application. Maintain a minimum of 14 days between applications. Use a NIS at 0.25% v/v or COC at 1% v/v. Refer to Table 1 at the end of this chapter for more specific information about grass control. Rain-free period is 1 hour. Fusilade also has some activity on bristly starbur (i.e. goathead or Texas sandspur).

PEANUT WEED CONTROL

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION/A	LBS ACTIVE INGREDIENT/A		
POST-EMERGENCE (continued)					
<i>chlorimuron</i> Classic 25DF	2	0.5 oz	0.008	-/ 45 D	Make one application per season as an over-the-top treatment for mid-season Florida beggarweed and bristly starbur control or suppression. Under favorable conditions—good soil moisture, moderate temperatures, and high relative humidity—other species such as cocklebur, ragweed, and sicklepod may be suppressed. Avoid applications during periods of drought/heat stress because of potential for poor weed control and crop injury. Applications of Classic may not provide acceptable control of Florida beggarweed that has escaped control or is re-growing after a previous application of Cadre. Include nonionic surfactant at 1 qt/100 gals spray solution with all Classic applications. Addition of ammonium sulfate (2 lb/A) or feed grade urea (2 gal/A) improves activity on bristly starbur. Classic can be applied from 60 days after peanut emergence to within 45 days of harvest. APPLICATIONS OF CLASSIC APPLIED FROM 60 DAYS AFTER CROP EMERGENCE TO 45 DAYS BEFORE HARVEST MAY CAUSE A SLIGHT INCREASE IN TSWV SYMPTOMS. Temporary yellowing of peanut foliage and reduction of canopy growth sometimes can occur. Can be tank-mixed with Bravo or 2,4-DB. However, combinations of Classic + 2,4-DB can result in significantly more foliar crop injury compared to Classic alone. Do not use on Spanish peanut. Do not tank-mix with elemental sulfur. Rain-free period for Classic is 1 hour. In recent weed-free trials conducted in Georgia, Classic has caused 7-11% yield reductions when applied to Georgia-06G and Tifguard. Significant yield losses have not occurred when Classic has been applied to Florida-07, Georgia Greener, and Georgia-07W. In 2013, GA-09B yields were reduced 5% when Classic was applied 74 DAE. Yields were not affected when applied to 60, 92, and 105 DAE.
NON-SELECTIVE APPLICATOR (NSA)					
<i>paraquat</i> 2SL 3SL	22	50% solution (2 lb ai/gal) 33% solution (3 lb ai/gal)		24 H/ 30 D	Section 24C Special Local Need label: For the salvage control/suppression of Palmer amaranth and Florida beggarweed. To prevent seed production in Palmer amaranth, apply within 2 weeks of pollen shed. Tractors should be operated at speeds of 5 MPH or less. NSAs that have performed well (> 85% control) in UGA tests include the following: GrassWorks Weed Wiper, Smucker's Top Crop Super Sponge, and LMC-Cross Wick-Bar. Do not apply more than 1 pt/A of a labeled <i>paraquat</i> formulation. In order for NSA's to be effective, at least 60-70% of the weed must be wicked/wiped. Additionally, this treatment may also be more effective on pigweed plants that are just starting to produce seed-heads. Rain-free time is 30 minutes. 24C Label must be in possession of user at the time of application. **All applicators of paraquat must complete the EPA's training program before mixing, loading and applying Refer to the following web-site for more information: https://www.epa.gov/pesticide-worker-safety/paraquat-dichloride-training-certified-applicators
HARVEST AID					
<i>carfentrazone</i> Aim 2EC	14	1-2 oz	0.156-0.031	12 H/ 7 D	Useful for the late-season desiccation/defoliation of annual morningglories (<i>Ipomoea</i> spp.). Aim is less effective on smallflower morningglory. Use in combination with either a NIS (0.25% v/v) or COC (1% v/v). Aim may cause peanut leaf spotting or burning. Use at least 15 GPA for optimum results. Do not graze or feed peanut hay to livestock. Only 1 application per season is permitted. Rain-free period is 6-8 hours.
<i>pyraflufen</i> ET 0.208EC	14	1-2 oz	0.0016-0.0032	12 H/ 7 D	Useful for the late-season desiccation/defoliation of annual morningglories (<i>Ipomoea</i> spp.). Use in combination with a NIS at 0.25% v/v (1 qt/100 gals). Apply a minimum of 7 days before harvest. Use at least 15 GPA for optimum results. ET will cause peanut leaf speckling/burn. Rain-free period is 1 hour.

PLANT GROWTH REGULATOR FOR RUNNER TYPE PEANUTS POST-EMERGENCE

Scott Monfort, Extension Peanut Agronomist

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION/A	LBS ACTIVE INGREDIENT/A		
<i>prohexadione calcium</i> Apogee or Kudos 27.5 WDG		3.63-5.44 oz	0.061 – 0.092	12 H/ 45 D	<p>To Suppress Excessive Vine Growth: Use the recommended reduced rate as the labeled rate (7.25 oz/Acre) is too excessive on runner type peanuts.</p> <p>The use of a growth regulator is only recommended on irrigated acres where vine growth is excessive. Use of a growth regulator in non-irrigated or in irrigated fields with reduced vine growth will lead to stunted growth and potential yield loss. The growth regulator is Not recommended on non-irrigated peanuts or in fields that are under stress due to lack of moisture, disease pressure, or other stress conditions.</p> <p>First application = when greater than 50% lateral vines are touching in the row middles (not at 50% lapped --- this will be too early). Second application = 14 to 21 days after 1st application. Apply in a minimum of 20 GPA of water.</p> <p>Always add a nitrogen source with Apogee or Kudos. Apply 1 lb of spray grade ammonium sulfate per acre or one pint of UAN (28%, 30% or 32% nitrogen solution) per acre. Add 1 quart/acre of Crop Oil Concentrate. May not need depending on tank mix partners. Rain fastness is 6-8 hours.</p>

WEED RESPONSE TO BURNDOWN HERBICIDES USED IN PEANUT

Eric P. Prostko and A. Stanley Culpepper, Extension Agronomists – Weed Science

WEED SPECIES	BURNDOWN TREATMENT ¹							
	<i>2,4-D</i> ³	<i>glyphosate acid</i> ²	<i>glyphosate acid</i> ² + <i>2,4-D</i> ³	<i>glyphosate acid</i> ² + Valor	<i>glyphosate acid</i> ² + Aim or ET	<i>paraquat</i>	<i>paraquat</i> + <i>2,4-D</i>	<i>paraquat</i> + Valor ⁴
GRASSES / SEDGES								
annual bluegrass	N	E	E	E	E	G-E	G-E	
bermudagrass	N	F	F	F	F	P	P	
crabgrass	N	E	G-E	E	E	G		
goosegrass	N	E	G-E	E	E	F-G		
Italian ryegrass	N	G-E	G	G	G	P-F	P-F	
johnsongrass	N	G-E	G	G-E	G-E	P		
little barley	N	E	E	E	E	G	G	
sandbur	N	E	G-E	E	E	G		
Texas panicum	N	E	G-E	E	E	G		
volunteer corn	N	E	E	E	E	F-G		
purple nutsedge	N	F-G	F-G	G	F-G	P-F		
yellow nutsedge	N	F	F	F	F	P-F		
BROADLEAVES								
bristly starbur	G	G-E	G-E	E	E	E		
buttercup	G	G-E	E	G-E	G-E	E		
Carolina geranium	F	P-F	G	G	F-G	G-E	G-E	
chickweed	P	E	E	E	E	E	E	
citronmelon	F	G-E	E	E	E	F		
cocklebur	E	E	E	E	E	G-E		
coffee senna	G	E	E	E	E	F		
corn spurry	P-F	G-E	G-E			F-G		
cowpea	G	E			E	E		
cudweed	P-F	G-E	G-E	E		F-G		
curly dock	P-F	F	F-G	F	F	P	P-F	
eveningprimrose	E	P-F	E	F-G	F	P-F	E	F-G

WEED RESPONSE TO BURNDOWN HERBICIDES USED IN PEANUT

WEED SPECIES	BURNDOWN TREATMENT ¹							
	2,4-D ³	glyphosate acid ²	glyphosate acid ² + 2,4-D ³	glyphosate acid ² + Valor	glyphosate acid ² + Aim or ET	paraquat	paraquat + 2,4-D	paraquat + Valor ⁴
BROADLEAVES (continued)								
eclipta	P	F			G-E	F		G
Florida beggarweed	P-F	E	E	E	E	E		
Florida pusley	F	F	G	F-G	G	F		G
field pansy	P-F	F-G	F-G	G		G-E		
hemp sesbania	G-E	P-F	E		G-E	F		
henbit	P-F	F	G-E	E	E	G	G-E	
horsenettle	F	F			P-F	P-F		
Horseweed	F-G	F-G	G-E	F-G	G	F	G	F
ALS-resistant	F-G	F-G	G-E	F-G	G	F	G	F
Glyphosate-resistant	F-G	P	F-G	P	F	F	G	F
lambsquarters	E	G	G		G-E	F-G		
morningglory, Ipomoea	G	F	E	E	E	F-G		
morningglory, smallflower	F	G	E	E	G-E	P		
Pennsylvania smartweed	F	G	G		G-E	P		
pigweed	G-E	G-E	E	E	E	G	G-E	G-E
ALS-resistant	G-E	G-E	G-E	E	E	G	G-E	G-E
Glyphosate-resistant	G-E	P	F-G	P	F	G	G-E	G-E
prickly sida	F-G	F-G	G		F-G	P-F		
purslane	G-E	F-G	G-E	G	F-G	G		
ragweed	E	G	E		G-E	G		
redweed	F	G			G-E	F		
shepherdspurse	G	G			G	G	G-E	
sicklepod	F-G	G-E	E	E	G-E	E		
speedwell	P-F	G-E	G-E	E	E	F	G	
spurred anoda	F-G	G			G	F-G		
swinecress	F-G	F-G	G	F-G	F-G	P	P-F	
tropic croton	F	G-E	G-E	E	G-E	F		

WEED RESPONSE TO BURNDOWN HERBICIDES USED IN PEANUT

WEED SPECIES	BURNDOWN TREATMENT ¹							
	2,4-D ³	glyphosate acid ²	glyphosate acid ² + 2,4-D ³	glyphosate acid ² + Valor	glyphosate acid ² + Aim or ET	paraquat	paraquat + 2,4-D	paraquat + Valor ⁴
BROADLEAVES (continued)								
tropical spiderwort	G-E	P	G-E	F	Aim = G-E ET = P-F	G	G-E	
velvetleaf	F-G	G			E	P		
vines (maypop, trumpet creeper, bigroot mg)	F	P-F			P-F	P		
Virginia pepperweed	G-E	G			G	P-F	G-E	
volunteer peanuts	P	F	F	F-G	F	P	P	F-G
volunteer sweet potato	G		G		F-G			
wild lettuce	G	G	G-E	E	G-E	P		
wild poinsettia	F-G	G			G-E	G-E		
wild radish	G	F-G	E	G	G	F	F-G	G
COVER CROPS								
clover	F	F	F-G		F	F-G		
lupine	G	G	G		G	F-G		
small grains	N	E	G-E	E	E	G		G
vetch	G	F	G-E	F	F	F		

Key:
 E – 90% or better control
 G – 80% to 90% control
 F – 60% to 80% control
 P – 30% to 60% control
 N – < 30% control

¹ Application rates per acre: 2,4-D, 1 pt; glyphosate acid, 0.75 lb a.e.; paraquat, 0.63 lb a.i.; Valor, 1 to 2.0 oz (Note: if 3 oz/A of Valor is used, burndown control may be better than indicated and residual control will be increased); Aim, 1-2 oz/A; ET, 0.5-2 oz/A.

² Mixing herbicides with glyphosate occasionally reduces grass control (including cover crops). This is more likely to occur with large weeds in dry conditions.

³ Labels for 2,4-D are ambiguous concerning the waiting period between application and planting.

⁴ Use a NIS (0.25% v/v) or COC (1% v/v) with this tank-mixture. A COC may be preferred if weeds are large

NOTE: Dicamba is not registered for pre-plant burndown use in peanut!

WEED RESPONSE TO HERBICIDES USED IN PEANUTS

Eric P. Prostko, Extension Agronomist – Weed Science

	PPI/PRE-EMERGENCE ^{1,2}					PRE-EMERGENCE				POST-EMERGENCE	
	Prowl Pendimax Sonalan	Dual Magnum ³	Propel Outlook	Pursuit	Strongarm	Warrant	Spartan Charge	Valor	Zidua ⁴	Paraquat ⁵	Paraquat + Storm
PERENNIALS											
bermudagrass	P	P	P	P	P	P		P	P	P	P
johnsongrass-rhizome	P	P	P	P	P	P		P	P	P	P
nutsedge, purple	P	P	P	G	P-F	P		P	P	P-F	F
nutsedge, yellow	P	F-G	F	F-G	P-F	P-F		P	P	P-F	F-G
broadleaf signalgrass	G-E	F-G	F	P	P	F-G		P	F-G	G	G
crabgrass	E	E	E	F	P	E		P	E	F-G	F-G
crowfootgrass	E	E	E	P	P	E		P	E	G	G
fall panicum	G	G	G	P-F	P	G		P	G	G	G
goosegrass	E	E	E	F	P	E		P	E	F-G	F-G
johnsongrass-seedling	E	F	F	G	P	F		P		G	G
sandbur	E	F-G	F-G		P	F-G		P	F-G	F	F-G
Texas panicum	G-E	P-F	P	P-F	P-F	P-F		P	P-F	G-E	G-E
BROADLEAVES											
bristly starbur	P	P	P	F	E	P		F		P-F	F-G
burgherkin	P	P	P	E	G	P		G		F	G
carpetweed	G	P-F	G	F-G	G	P				F-G	G
citronmelon	P	P	P	P	G	P		G		F	G
cocklebur	P	P	P	G-E	G-E	P		P		G	G-E
coffee senna	P	P	P	F-G	P	P		P-F		F	E
copperleaf, hophornbeam	P	P	F	P	G-E	P		G-E		P	G
cowpea	P	P	P	P	P	P		P-F		F-G	F

WEED RESPONSE TO HERBICIDES USED IN PEANUTS

	PPI/PRE-EMERGENCE ^{1,2}					PRE-EMERGENCE				POST-EMERGENCE	
	Prowl Pendimax Sonalan	Dual Magnum ³	Propel Outlook	Pursuit	Strongarm	Warrant	Spartan Charge	Valor	Zidua ⁴	Paraquat ⁵	Paraquat + Storm
BROADLEAVES (continued)											
crotalaria, showy	P	P	P			P		G			F-G
croton, tropic	P	P	P	P	F-G	P		G		P	G
dayflower, Benghal tropical spiderwort	P	G-E	F	G	G	G-E		F	G-E	G	G
eclipta	P	P-F	P-F	P	G-E			G-E		P-F	F-G
Florida beggarweed	P	P-F	P-F	P	F-G	P-F		G-E	F	G-E	G-E
Florida pusley	E	G-E	G-E	G	G-E	G		G-E		P	P
groundcherry, cutleaf	P	G	G								
jimsonweed	P			G	G-E			G		P	F
hairy indigo	P	F			G			G		F	
hemp sesbania	P	P	P	P	P-F	P		G-E	F-G		G
horseweed						P		G-E		P	P
lambsquarters	E	F	G	F	G-E	F		G-E		F	F-G
morningglory spp.	P	P	P	G	F-G	P		F-G		P	F
cypressvine	P	P	P	G		P		G		F-G	F-G
entireleaf/ivyleaf	P	P	P	G	F-G	P		F-G		F	G
pitted	P	P	P	G	F-G	P		F		F	G
purple moonflower	P	P	P			P				F	G
red	P	P	P	G	F	P		G		F	G
smallflower	P	P	P	E	G	P		G-E		P	G-E
tall	P	P	P	G		P		F-G		F	G
pigweed	G	G	G	E	G	G	G-E	E	G	F	G-E
ALS-resistant	G	G	G	P	P	G	G-E	E	G	F	G-E
poorjoe											

WEED RESPONSE TO HERBICIDES USED IN PEANUTS

	PPI/PRE-EMERGENCE ^{1,2}					PRE-EMERGENCE				POST-EMERGENCE	
	Prowl Pendimax Sonalan	Dual Magnum ³	Propel Outlook	Pursuit	Strongarm	Warrant	Spartan Charge	Valor	Zidua ⁴	Paraquat ⁵	Paraquat + Storm
BROADLEAVES (continued)											
prickly sida	P	F	F	G-E	F-G	F		G-E		F	G
primrose, cutleaf evening										P	G (+2,4-DB)
purslane	G-E	G	G	G		G		G-E		G	G
ragweed, common	P	P	F-G	P	G-E			G-E		P-F	G
redweed	P				G			G-E		F	G
spurred anoda	P	P	P		F-G			F		P	G
sicklepod	P	P	P	P	P	P		P		G-E	G-E
smartweed	P			G	G			P-F		G-E	G
spider flower	P	P	P	G							
spurge spp.	P	P-F	P-F					G-E			
velvetleaf	P	P	P	P-F	G-E			F		F	F-G
wild poinsettia	P	P	P	E	G-E			P-F		F	G
wild radish	P	P	P	E		P			P-F	F	G

Abbreviations:
 E – Excellent (> 90%)
 G – Good (80-89%)
 F – Fair (70-79%)
 P – Poor (< 70%)
 If no letter is given, response is unknown.
 PPI = Pre-plant Incorporated,
 PRE = Pre-emergence.

¹ Ratings for Pursuit PPI and PRE are similar.

² Ratings for Dual, Lasso, and Outlook/Propel PRE and AC are similar. See remarks for additional information.

³ The generic formulations of *metolachlor* (Parallel PCS, Stalwart, Me-Too-Lachlor) have not provided the same length of residual control of certain weeds as similar rates of Dual Magnum formulations in some UGA field trials.

⁴ Zidua should not be applied PRE in peanut due to potential injury concerns. Zidua provides residual control of certain weeds when tank-mixed with POST herbicides such as Cadre, Cobra, or Ultra Blazer.

⁵ Commercially available as Helmqat, Firestorm or Parazone or Gramoxone SL.

WEED RESPONSE TO HERBICIDES USED IN PEANUTS

	POST-EMERGENCE												
	Strongarm**	Paraquat + Basagran	2,4-DB	Pursuit	Basagran	Ultra Blazer	Cobra	Storm	Cadre	Fusilade	Select	Poast	Classic
PERENNIALS													
bermudagrass	P	P	P	P	P	P	P	P	P	G	G	F-G	P
johnsongrass-(rhizome)	P	P	P	P	P	P	P	P	F-G	G-E	G	F-G	P
nutsedge, purple		F	P	G	P	P	P	P	G-E	P	P	P	P
nutsedge, yellow		F-G	P	F-G	G	P	P-F	F	G-E	P	P	P	F-G
GRASSES													
broadleaf signalgrass	P	G	P	P	P	P	P-F	P	G	G	G-E	G-E	P
crabgrass	P	F-G	P	P-F	P	P	P-F	P	G	G	G-E	G-E	P
crowfootgrass	P	G	P	P-F	P	P	P	P	G	F-G	G	F-G	P
fall panicum	P	G	P	P	P	F	P	P	G	G-E	G-E	G-E	P
goosegrass	P	F-G	P	P	P	P	P	P	F	G	G	G	P
johnsongrass-seedling	P	G	P	F	P	P	P	P	F-G	G-E	G-E	G-E	P
sandbur	P	F-G	P		P	P	P-F	P	G	G	G	G	P
Texas panicum	P	G-E	P	P-F	P	P	P	P	F-G	G	G-E	G-E	P
BROADLEAVES													
bristly starbur	E	F	P-F	P-F	G	P-F	G	F-G	F	F	P	P	F
burgherkin		F	F	F	P	G	G	F	G-E	P	P	P	P
carpetweed		P	P	F-G	P	G-E	G-E	G	F-G	P	P	P	
citronmelon		F	G	P	P	F	G	F	G	P	P	P	P
cocklebur	E	G	E	E	E	G	G-E	E	E	P	P	P	F
coffee senna		E	F-G	F	G	P	P-F	F	G	P	P	P	P
copperleaf, hophornbeam	P	P	P	P	P	G-E	G-E	F	P-F	P	P	P	P
cowpea		P-F	P-F	P	P	P-F	P-F	P-F	P-F	P	P	P	F-G
crotalaria, showy					P	E	E	G-E		P	P	P	

WEED RESPONSE TO HERBICIDES USED IN PEANUTS

	POST-EMERGENCE												
	Strongarm**	Paraquat + Basagran	2,4-DB	Pursuit	Basagran	Ultra Blazer	Cobra	Storm	Cadre	Fusilade	Select	Poast	Classic
BROADLEAVES (continued)													
croton, tropic	P	P	P	P	P	E	E	G-E	P	P	P	P	P
dayflower, Benghal tropical spiderwort	G	G	P	F-G	G	P	P	P	F-G	P	P	P	F
eclipta	G-E	F	P	P	G	F	F	G	P-F	P	P	P	P
Florida beggarweed	P-F	G-E	P	P	P	P	P-F	P	F-G	P	P	P	F-G
Florida pusley		P	P	P	P	P	F-G	P	P	P	P	P	P
groundcherry, cutleaf		F-G			P	G	G	F-G		P	P	P	
jimsonweed		E	P	F-G	E	E	E	G	E	P	P	P	
hairy indigo			F	P	P	G	G	F	F	P	P	P	F-G
hemp sesbania				P	P	E	E	G-E	P	P	P	P	F-G
horseweed ALS-resistant	G P	P	P	P	P	P	P	P	P	P	P	P	F P
lambsquarters		F	F	P	F	P-F	P-F	F	P-F	P	P	P	P
morningglory spp.	G-E	F-G	F-G	G	F	G-E	G-E	G	G	P	P	P	
cypressvine		G-E	F	G	G	G	G-E	G	G	P	P	P	
entireleaf/ivyleaf	G-E		G	F-G	P	G	F-G	F	G	P	P	P	
pitted	G-E		F-G	G	P	G-E	G	F-G	G	P	P	P	
purple moonflower	F-G		F-G	P	P	G-E	G-E	G	F	P	P	P	P
red			G		F-G	G-E	G-E	G-E		P	P	P	
smallflower	G-E	G-E	F	E	E	G-E	G-E	G-E	E	P	P	P	
tall			G		P	G	G	F-G	G	P	P	P	
pigweed ALS-resistant	P P	F-G F-G	F F	E P	P P	G-E G-E	G-E G-E	G G	E P	P P	P P	P P	F-G P

WEED RESPONSE TO HERBICIDES USED IN PEANUTS

	POST-EMERGENCE												
	Strongarm**	Paraquat + Basagran	2,4-DB	Pursuit	Basagran	Ultra Blazer	Cobra	Storm	Cadre	Fusilade	Select	Poast	Classic
BROADLEAVES (continued)													
poorjoe			F			G	G			P			
prickly sida		G	P	P-F	G	P	G	G	G	P	P	P	P
primrose, cutleaf evening	P	F (+2,4-DB)	F	P	P	P	P	P	P	P	P	P	P
purslane		G	G	P-F	G	E	E	G-E	P-F	P	P	P	
ragweed, common	E	F	F	P	F	E	E	G	F	P	P	P	P-F
redweed		G	P	P	G	P	F	G	G	P	P	P	P
spurred anoda		F-G	P		G	P	P	F	G	P	P	P	
sicklepod	P	G	F	P	P	P	P-F	P	G-E	P	P	P	P-F
smartweed		G	P	G-E	G-E	G-E	G-E	G-E	F-G	P	P	P	P
spider flower				F-G		G	G	F	F-G	P	P	P	F
spurge spp.			P	P	P	F	F	F		P	P	P	P
velvetleaf		G	P	P-F	G	P-F	G	F-G		P	P	P	
wild poinsettia	P-F	G-E	P	P-F	P	G-E	G-E	G	E	P	P	P	P
wild radish	G	F	P	G-E	P-F	E	E	G	E	P	P	P	P

<p>Abbreviations: E – Excellent (> 90%) G – Good (80-89%) F – Fair (70-79%) P – Poor (< 70%)</p>	<p>If no symbol is given, response is unknown. ** 24(c) label for use in Georgia only for tropical spiderwort.</p>
--	---

SUMMARY OF PEANUT GRASS HERBICIDES

	HERBICIDE					
	Fusilade DX	Poast	Poast Plus	Select/Arrow/Others (2 lb ai/gal)	Select Max/TapOut (0.97 lb ai/gal)	Section Three Shadow (3 lb ai/gal)
Maximum Rate/A/ Season	48 oz	40 oz	60 oz	32 oz	64 oz	21.33 oz
Maximum Rate/A/ Application	24 oz	24 oz	24 oz	16 oz	32 oz	10.67 oz
broadleaf signalgrass	12 oz (2-4")	16 oz (up to 8")	24 oz (up to 8")	6-8 oz (2-6")	9-16 oz (2-6")	3.33 oz
crabgrass	12 oz (1-2")	16 oz (up to 6")	24 oz (up to 6")	6-8 oz (2-6")	9-16 oz (2-6")	3.33 oz (1-4")
crowfootgrass	NL*	NL	NL	6-8 oz (2-6")	9-16 oz (2-6")	NL
field sandbur	12 oz (2-4")	20 oz (up to 3")	30 oz (up to 3")	6-8 oz (2-6")	9-16 oz (2-6")	NL
goosegrass	8 oz (2-4")	16 oz (up to 6")	24 oz (up to 6")	6-8 oz	9-16 oz (2-6")	NL
Texas panicum	12 oz (2-8")	16 oz (up to 8")	24 oz (up to 8")	6-8 oz (2-6")	9-16 oz (2-6")	3.33 oz (1-4")
rhizome johnsongrass	12-24 oz (1 st) (8-18")	24 oz (1 st) (up to 25")	36 oz (1 st) (up to 25")	8-16 oz (1 st) (12-24")	12-32 oz (1 st) (12-24")	5.33-10.67 oz (1 st) (12-24")
	8-24 oz (2 nd) (6-12")	16 oz (2 nd) (up to 12")	24 oz (2 nd) (up to 12")	6-8 oz (2 nd) (6-18")	9-24 oz (2 nd) (6-18")	4-5.33 oz (2 nd) (6-18")
bermudagrass	12-24 oz (1 st) (4-8" runners)	24 oz (1 st) (up to 6" stolon)	36 oz (1 st) (up to 6" stolon)	8-16 oz (1 st) (3-6" runners)	12-32 oz (1 st) (3-6" runners)	5.33-10.67 oz (1 st) (up to 6" runners)
	8-24 oz (2 nd) (4-8" runners)	16 oz (2 nd) (up to 4" stolon)	24 oz (2 nd) (up to 4" stolon)	8-16 oz (2 nd) (3-6" runners)	12-32 oz (2 nd) (3-6" runners)	5.33-10.67 oz (up to 6" runners)
volunteer corn	6 oz (12-24" tall)	16 oz (≤ 20" tall)	24 oz (≤ 20" tall)	4-6 oz (4-12" tall)	≤ 12" = 6-12 oz 13-24" = 9-14 oz 25-36" = 12-16 oz	≤ 12" = 2.67-4 oz 13-24" = 4-5.33 oz 25-36" = 5.33-6.67 oz

* NL= crowfootgrass was not listed on the product label.

SUGGESTED HERBICIDE PROGRAMS FOR THE CONTROL OF TROPICAL SPIDERWORT (BENGHAL DAYFLOWER) IN PEANUT***:

PROGRAM 1

- a) **PRE Immediately After Planting:** Valor at 3 oz/A + Dual Magnum 7.62EC or generic *metolachlor* (Stalwart, Parallel PCS, Me-Too-Lachlor) at 1 pt/A or Warrant 3ME at 3 pt/A **and**
- b) **POST when spiderwort is 1-2” tall:** Cadre/Impose 2AS at 4 oz/A or Strongarm 84WG at 0.45 oz/A + Dual Magnum 7.62EC or generic *metolachlor* (Stalwart, Parallel PCS, Me-Too-Lachlor) at 1 pt/A or Warrant 3ME at 3 pt/A or Zidua 4.17SC @ 2.45 oz/A.
 - * At plant applications of Dual or Warrant are more effective for TSW/BD control at later peanut planting dates (May 15 or later) because the majority of emergence occurs after June 1.

PROGRAM 2

- a) **AT-CRACK (within 28 days after peanut cracking):** Apply paraquat 2SL formulations at 12 oz/A A or paraquat 3SL formulations at 8 oz/A Storm 4EC at 16 oz/A or Basagran 4EC @ 8 oz/A + Dual Magnum 7.62EC or generic *metolachlor* (Stalwart, Parallel PCS, Me-Too-Lachlor) at 1 pt/A or Warrant 3ME at 3 pt/A or Zidua 4.17SC @ 2.45 oz/A **and**
- b) **POST (~2-3 weeks after at-crack spray):** Apply Cadre/Impose 2AS at 4 oz/A or Strongarm 84WG at 0.45 oz/A + Dual Magnum 7.62EC or generic *metolachlor* (Stalwart, Parallel PCS, Me-Too-Lachlor) at 1 pt/A or Warrant 3ME at 3 pt/A or Zidua 4.17SC @ 2.45 oz/A.

* When using Dual Magnum or generic *metolachlor* POST in combination with Cadre/Impose, *paraquat*, or Strongarm, additional spray adjuvants (NIS, COC) are not necessary.

The maximum amount/A/year of Dual Magnum that can be applied is 2.8 pt/A. The maximum amount/A/year of Stalwart, Parallel PCS, or Me-To-Lachlor that can be applied is 2.66 pt/A.

The maximum amount of Warrant that can be applied PRE + POST is 6 pt/A/year.

Maximum amount of Zidua that can be applied is 4 oz/A/year of 85WG or 6.5 oz/A/year of 4.17SC.

When Warrant or Zidua is applied POST, a NIS (0.25% v/v) is needed.

** **Zidua should not be applied prior to peanut emergence due to potential injury concerns.**

*****Twin rows and conventional tillage (plowing) are also useful for the management of tropical spiderwort/Benghal dayflower.**

HERBICIDE PROGRAMS FOR PEANUT IN GEORGIA - 2020

SYSTEM	TILLAGE	TIMING				
		PRE-PLANT BURNDOWN ^a	PPI	PRE	EPOST (~10-20 DAP ^b)	POST (~30-45 DAP)
non-irrigated (dryland)	strip-till ^c	<i>Glyphosate</i> or <i>Paraquat</i> + 2,4-D		No Rain in 7-10 DAP <i>Paraquat</i> + Prowl	<i>Paraquat</i> + Storm + Dual Magnum or Warrant or Zidua	ALS Resistance: Cobra or Ultra Blazer + Dual Magnum or Warrant or Zidua + 2,4-DB No ALS Resistance: Cadre ^d + (Dual Magnum or Warrant or Zidua) + 2,4-DB **A 4-way tank-mixture can be used if required (Cadre + Cobra or Ultra Blazer + 2,4-DB + Dual Magnum or Warrant or Zidua)
				Rain in 7-10 DAP <i>Paraquat</i> + Prowl + Valor + Strongarm ^d		
	conventional	Prowl or Sonalan	No PRE if rain is not expected in 7-10 DAP	<i>Paraquat</i> + Storm + Dual Magnum or Warrant or Zidua		
			Rain in 7-10 DAP Valor + Strongarm ^d			
irrigated	strip-till ^c	<i>Glyphosate</i> or <i>Paraquat</i> + 2,4-D		<i>Paraquat</i> + Prowl + Valor + Strongarm ^d		
	conventional			Prowl or Sonalan + Valor + Strongarm ^d		

^a Apply at least 7 days before planting. If there will be a long delay between the burndown application and planting (> 10 days), consider adding a residual pigweed herbicide (Valor or Dual Magnum or Warrant) to the burndown treatment.

^b DAP = days after planting.

^c Annual grass control in strip-tillage systems is often more difficult thus additional applications of a post-emergence grass herbicide (i.e. Fusilade, Poast, and Select) may be needed.

^d Before using Cadre and/or Strongarm, rotational crop restrictions must be considered.

** **SPECIAL NOTE:** Dual Magnum and Warrant are in the same herbicide family (*chloroacetamide*) and have the same mode of action (inhibit very long chain fatty acid synthesis, WSSA MOA #15). Zidua is not in the same herbicide family (*isoxazoline*) but has the same mode of action. Multiple applications (> 2) of these herbicides in a single year should be avoided to prevent or delay the evolution of resistance. These herbicides have no post-emergence activity.

